

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
20 September 2001 (20.09.2001)

PCT

(10) International Publication Number  
WO 01/68603 A2

(51) International Patent Classification<sup>7</sup>: C07D 209/52,  
A61K 31/403, A61P 3/04, 3/06, 3/10

David, A. [US/US]; 3 Easton Court, Lawrenceville, NJ  
08648 (US).

(21) International Application Number: PCT/US01/07151

(74) Agents: ALGIERI, Aldo, A. et al.; Bristol-Myers Squibb  
Co., P.O. Box 4000, Lawrenceville-Princeton Rd., Prince-  
ton, NJ 08543 (US).

(22) International Filing Date: 5 March 2001 (05.03.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/188,555 10 March 2000 (10.03.2000) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE,  
DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU,  
ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,  
LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ,  
PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT,  
TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(71) Applicant (*for all designated States except US*): BRIS-  
TOL-MYERS SQUIBB CO. [US/US]; P.O. Box 4000,  
Lawrenceville-Princeton Road, Princeton, NJ 08543 (US).

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): ROBL, Jeffrey,  
A. [US/US]; 7 Tulip Drive, Newtown, PA 18940 (US).  
SULSKY, Richard, B. [US/US]; 2 Coventry Square, West  
Trenton, NJ 08628 (US). AUGERI, David, J. [US/US];  
107 Carter Road, Princeton, NJ 08540 (US). MAGNIN,  
David, R. [US/US]; 40 Cottage Court, Hamilton, NJ 08690  
(US). HAMANN, Lawrence, G. [US/US]; 24 East Riding  
Drive, Cherry Hill, NJ 08003 (US). BETEBENNER,


Published:

— without international search report and to be republished  
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

WO 01/68603 A2

(54) Title: CYCLOPROPYL-FUSED PYRROLIDINE-BASED INHIBITORS OF DIPEPTIDYL PEPTIDASE IV AND METHOD



(57) Abstract: Dipeptidyl peptidase IV (DP 4) inhibiting compounds are provided having the formula (I) where x is 0 or 1 and y is 0 or 1 (provided that x = 1 when y = 0 and x = 0 when y = 1); n is 0 or 1; X is H or CN; and wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are as described herein. A method is also provided for treating diabetes and related diseases, especially Type II diabetes, and other diseases as set out herein, employing such DP 4 inhibitor or a combination of such DP 4 inhibitor and one or more of another antidiabetic agent such as metformin, glyburide, troglitazone, pioglitazone,

rosiglitazone and/or insulin and/or one or more of a hypolipidemic agent and/or anti-obesity agent and/or other therapeutic agent.

CYCLOPROPYL-FUSED PYRROLIDINE-BASED INHIBITORS OF  
DIPEPTIDYL PEPTIDASE IV AND METHOD

5           The present invention relates to cyclopropyl-fused  
pyrrolidine-based inhibitors of dipeptidyl peptidase IV  
(DP-4), and to a method for treating diabetes, especially  
Type II diabetes, as well as hyperglycemia, Syndrome X,  
diabetic complications, hyperinsulinemia, obesity,  
10   atherosclerosis and related diseases, as well as various  
immunomodulatory diseases and chronic inflammatory bowel  
disease, employing such cyclopropyl-fused pyrrolidines  
alone or in combination with another type antidiabetic  
agent and/or other type therapeutic agent.

15           Depeptidyl peptidase IV (DP-4) is a membrane bound  
non-classical serine aminodipeptidase which is located in  
a variety of tissues (intestine, liver, lung, kidney) as  
well as on circulating T-lymphocytes (where the enzyme is  
20   known as CD-26). It is responsible for the metabolic  
cleavage of certain endogenous peptides (GLP-1(7-36),  
glucagon) in vivo and has demonstrated proteolytic  
activity against a variety of other peptides (GHRH, NPY,  
GLP-2, VIP) in vitro.


25           GLP-1(7-36) is a 29 amino-acid peptide derived by  
post-translational processing of proglucagon in the small  
intestine. GLP-1(7-36) has multiple actions in vivo  
including the stimulation of insulin secretion,  
30   inhibition of glucagon secretion, the promotion of  
satiety, and the slowing of gastric emptying. Based on  
its physiological profile, the actions of GLP-1(7-36) are  
expected to be beneficial in the prevention and treatment  
of type II diabetes and potentially obesity. To support

this claim, exogenous administration of GLP-1(7-36) (continuous infusion) in diabetic patients has demonstrated efficacy in this patient population. Unfortunately GLP-1(7-36) is degraded rapidly in vivo and  
 5 has been shown to have a short half-life in vivo ( $t_{1/2} \approx 1.5$  min). Based on a study of genetically bred DP-4 KO mice and on in vivo/in vitro studies with selective DP-4 inhibitors, DP-4 has been shown to be the primary degrading enzyme of GLP-1(7-36) in vivo. GLP-1(7-36) is  
 10 degraded by DP-4 efficiently to GLP-1(9-36), which has been speculated to act as a physiological antagonist to GLP-1(7-36). Thus, inhibition of DP-4 in vivo should potentiate endogenous levels of GLP-1(7-36) and attenuate formation of its antagonist GLP-1(9-36) and thus serve to  
 15 ameliorate the diabetic condition.

In accordance with the present invention, cyclopropyl-fused pyrrolidine-based compounds are provided which inhibit DP-4 and have the structure

20

I



wherein x is 0 or 1 and y is 0 or 1 (provided that

25 x = 1 when y = 0 and

x = 0 when y = 1);

n is 0 or 1;


X is H or CN (that is cyano);

30 R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are the same or different and are independently selected from H, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, bicycloalkyl, tricycloalkyl, alkylcycloalkyl, hydroxyalkyl, hydroxyalkylcycloalkyl, hydroxycycloalkyl, hydroxybicycloalkyl, hydroxytricycloalkyl, bicycloalkylalkyl, alkylthioalkyl,


arylalkylthioalkyl, cycloalkenyl, aryl, aralkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl and cycloheteroalkylalkyl, all optionally substituted through available carbon atoms with 1, 2, 3, 4 or 5 groups selected from hydrogen, halo, alkyl, polyhaloalkyl, alkoxy, haloalkoxy, polyhaloalkoxy, alkoxycarbonyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, polycycloalkyl, heteroarylamino, arylamino, cycloheteroalkyl, cycloheteroalkylalkyl, hydroxy, hydroxyalkyl, nitro, cyano, amino, substituted amino, alkylamino, dialkylamino, thiol, alkylthio, alkylcarbonyl, acyl, alkoxycarbonyl, aminocarbonyl, alkynylaminocarbonyl, alkylaminocarbonyl, alkenylaminocarbonyl, alkylcarbonyloxy, alkylcarbonylamino, arylcarbonylamino, alkylsulfonylamino, alkylaminocarbonylamino, alkoxycarbonylamino, alkylsulfonyl, aminosulfonyl, alkylsulfinyl, sulfonamido or sulfonyl;

and  $R^1$  and  $R^3$  may optionally be taken together to form  $-(CR^5R^6)_m-$  where  $m$  is 2 to 6, and  $R^5$  and  $R^6$  are the same or different and are independently selected from hydroxy, alkoxy, cyano, H, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, cycloalkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, halo, amino, substituted amino, cycloheteroalkylalkyl, alkylcarbonylamino, arylcarbonylamino, alkoxycarbonylamino, aryloxy carbonylamino, alkoxycarbonyl, aryloxy carbonyl, or alkylaminocarbonylamino, or  $R^1$  and  $R^4$  may optionally be taken together to form  $-(CR^7R^8)_p-$  where  $p$  is 2 to 6, and  $R^7$  and  $R^8$  are the same or different and are independently selected from hydroxy, alkoxy, cyano, H, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, cycloalkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, halo, amino, substituted amino, cycloheteroalkylalkyl, alkylcarbonylamino, arylcarbonylamino, alkoxycarbonylamino, aryloxy carbonylamino,

alkoxycarbonyl, aryloxycarbonyl, or  
alkylaminocarbonylamino, or optionally R<sup>1</sup> and R<sup>3</sup> together



with R<sup>4</sup> form a 5 to 7 membered ring containing a  
total of 2 to 4 heteroatoms selected from N, O, S, SO, or  
5 SO<sub>2</sub>;




or optionally R<sup>1</sup> and R<sup>3</sup> together with form  
a 4 to 8 membered cycloheteroalkyl ring wherein the  
cycloheteroalkyl ring has an optional aryl ring fused  
thereto or an optional 3 to 7 membered cycloalkyl ring  
10 fused thereto;


and including pharmaceutically acceptable salts  
thereof, and prodrug esters thereof, and all  
stereoisomers thereof.

Thus, the compounds of formula I of the invention  
15 include the following structures

IA



IB



20

In addition, in accordance with the present  
invention, a method is provided for treating diabetes,  
especially Type II diabetes, as well as impaired glucose  
homeostasis, impaired glucose tolerance, infertility,  
25 polycystic ovary syndrome, growth disorders,  
frailty, arthritis, allograft rejection in transplant-  
ation, autoimmune diseases (such as scleroderma and

multiple sclerosis), various immunomodulatory diseases (such as lupus erythematosus or psoriasis), AIDS, intestinal diseases (such as necrotizing enteritis, microvillus inclusion disease or celiac disease),  
5 inflammatory bowel syndrome, chemotherapy-induced intestinal mucosal atrophy or injury, anorexia nervosa, osteoporosis, Syndrome X, dysmetabolic syndrome, diabetic complications, hyperinsulinemia, obesity, atherosclerosis and related diseases, as well as inflammatory bowel  
10 disease (such as Crohn's disease and ulcerative colitis), wherein a therapeutically effective amount of a compound of structure I (which inhibits DP 4) is administered to a human patient in need of treatment.

The conditions, diseases, and maladies collectively  
15 referenced to as "Syndrome X" or Metabolic Syndrome are detailed in Johannsson *J. Clin. Endocrinol. Metab.*, 82, 727-734 (1997).

In addition, in accordance with the present invention, a method is provided for treating diabetes and  
20 related diseases as defined above and hereinafter as well as any of the other disease states mentioned above, wherein a therapeutically effective amount of a combination of a compound of structure I and one, two, three or more of other types of antidiabetic agent(s)  
25 (which may be employed to treat diabetes and related diseases) and/or one, two or three or more other types of therapeutic agent(s) is administered to a human patient in need of treatment.

The term "diabetes and related diseases" refers to  
30 Type II diabetes, Type I diabetes, impaired glucose tolerance, obesity, hyperglycemia, Syndrome X, dysmetabolic syndrome, diabetic complications, dysmetabolic syndrome, and hyperinsulinemia.

The conditions, diseases and maladies collectively  
35 referred to as "diabetic complications" include retinopathy, neuropathy and nephropathy, and other known complications of diabetes.

The term "other type(s) of therapeutic agents" as employed herein refers to one or more antidiabetic agents (other than DP4 inhibitors of formula I), one or more anti-obesity agents, and/or one or more lipid-modulating agents (including anti-atherosclerosis agents), and/or one or more infertility agents, one or more agents for treating polycystic ovary syndrome, one or more agents for treating growth disorders, one or more agents for treating frailty, one or more agents for treating arthritis, one or more agents for preventing allograft rejection in transplantation, one or more agents for treating autoimmune diseases, one or more anti-AIDS agents, one or more anti-osteoporosis agents, one or more agents for treating immunomodulatory diseases, one or more agents for treating chronic inflammatory bowel disease or syndrome and/or one or more agents for treating anorexia nervosa.



The term "lipid-modulating" agent as employed herein refers to agents which lower LDL and/or raise HDL and/or lower triglycerides and/or lower total cholesterol and/or other known mechanisms for therapeutically treating lipid disorders.

In the above methods of the invention, the compound of structure I will be employed in a weight ratio to the antidiabetic agent or other type therapeutic agent (depending upon its mode of operation) within the range from about 0.01:1 to about 500:1, preferably from about 0.1:1 to about 100:1, more preferably from about 0.2:1 to about 10:1.


Preferred are compounds of formula I wherein R<sup>3</sup> is H or alkyl, R<sup>1</sup> is H, alkyl, cycloalkyl, bicycloalkyl, tricycloalkyl, alkylcycloalkyl, hydroxyalkyl, hydroxytricycloalkyl, hydroxycycloalkyl, hydroxybicycloalkyl, or hydroxyalkylcycloalkyl, R<sup>2</sup> is H or alkyl, n is 0, X is CN, x is 0 or 1 and y is 0 or 1.





Most preferred are preferred compounds of formula

I as described above where X is CN or CN ,

and/or wherein the fused cyclopropyl group is


identified as  .

Thus, preferred compounds of formula I of the invention will include the moiety:




10

or



Particularly preferred are the following compounds:


15



[1S, 2(2S), 3S, 5S]


wherein R<sup>1</sup> is alkyl, cycloalkyl, bicycloalkyl, tricycloalkyl, alkylcycloalkyl, hydroxyalkyl, hydroxycycloalkyl, hydroxyalkylcycloalkyl, hydroxybicycloalkyl or hydroxytricycloalkyl;

20



[1R,2S,3(2S),5S]

wherein R<sup>1</sup> is alkyl, cycloalkyl, bicycloalkyl, tricycloalkyl, hydroxybicycloalkyl, hydroxytricycloalkyl, alkylcycloalkyl, hydroxyalkyl, hydroxycycloalkyl or hydroxyalkylcycloalkyl as well as the following:



10

and

15 Compounds of the structure I may be generated by the methods as shown in the following reaction schemes and the description thereof.

Referring to Reaction Scheme 1, compound 1, where PG<sub>1</sub> is a common amine protecting group such as Boc, Cbz, or Fmoc and X<sup>1</sup> is H or CO<sub>2</sub>R<sup>9</sup> as set out below, may be generated by methods as described herein or in the literature (for example see Sagnard et al, Tet-Lett., 1995, 36, pp. 3148-3152, Tverezovsky et al, Tetrahedron,

20

1997, 53, pp. 14773-14792, Hanessian et al, Bioorg. Med. Chem. Lett., 1998, 8, p. 2123-2128). Removal of the PG<sub>1</sub> group by conventional methods (e.g. (1) TFA or HCl when PG<sub>1</sub> is Boc, or (2) H<sub>2</sub>/Pd/C, TMSI when PG<sub>1</sub> is Cbz, or (3) Et<sub>2</sub>NH when PG<sub>1</sub> is (Fmoc) affords the free amine 2. Amine 2 may be coupled to various protected amino acids such as 3 (where PG<sub>2</sub> can be any of the PG<sub>1</sub> protecting groups) using standard peptide coupling conditions (e.g. EDAC/HOAT, i-BuOCOC1/TEA, PyBop/NMM) to afford the corresponding dipeptide 4. Removal of the amine protecting group PG<sub>2</sub> provides compound Ia of the invention where X=H.


In the case where X<sup>1</sup>=CO<sub>2</sub>R<sup>9</sup> (where R<sup>9</sup> is alkyl or aralkyl groups such as methyl, ethyl, t-butyl, or benzyl), the ester may be hydrolyzed under a variety of conditions, for example with aqueous NaOH in a suitable solvent such as methanol, THF, or dioxane, to provide the acid 5. Conversion of the acid group to the primary carboxamide, affording 6, may be effected by activation of the acid group (e.g. employing i-BuOCOC1/TEA or EDAC) followed by treatment with NH<sub>3</sub> or an ammonia equivalent in a solvent such as dioxane, ether, or methanol. The amide functionality may be converted to the nitrile group by a variety of standard conditions (e.g. POCl<sub>3</sub>/pyridine/imidazole or cyanuric chloride/DMF or trifluoroacetic anhydride, THF, pyridine) to give 7. Finally, removal of the PG<sub>2</sub> protecting group similar to above provides compound of the invention Ib.

In a different sequence (Scheme 2), compound 1 where X<sup>1</sup> is CO<sub>2</sub>R<sup>9</sup> may be saponified to the acid and subsequently amidated as described above to give amide 8. Removal of the PG<sub>1</sub> group followed by peptide coupling to 3 affords compound 6, an intermediate in the synthesis of Ib.

Alternately, the carboxamide group in 8 may be converted to the nitrile as described above to give compound 9. Deprotection of PG<sub>1</sub> affords 10 which may be


subject to standard peptide coupling conditions to afford 7, an intermediate in the synthesis of Ib. Compound 10 may also be generated by oxidation of the amine 2 (e.g. NCS) followed by hydrolysis and subsequent cyanide treatment. Compound 10 may be obtained as a mixture of stereoisomers or a single isomer/diastereomer which may be epimerized (employing conventional procedures) to afford a mixture of stereoisomers.

## 10 Scheme 1




a.  $\text{PG}_1 = \text{Boc}$ , TFA or HCl;  $\text{PG}_1 = \text{Cbz}$ ,  $\text{H}_2/\text{Pd/C}$  or TMSI;  $\text{PG}_1 = \text{Fmoc}$ ,  $\text{Et}_3\text{NH}$  b. EDAC, HOBT, DMF or  $i\text{-BuOCOCi/TEA}$  or PyBop, NMM c.  $\text{PG}_2 = \text{PG}_1$ , (see conditions for a) d. LiOH or NaOH MeOH or THF/ $\text{H}_2\text{O}$  or dioxane e.  $i\text{-BuOCOCi/ NMM}$  or  $i\text{-BuOCOCi/TEA}$  or EDAC, then  $\text{NH}_3$  in dioxane or  $\text{Et}_2\text{O}$  f.  $\text{POCl}_3$ , pyridine, imidazole or cyanuric chloride, DMF or TFAA, THF, pyridine.

Scheme 2



5


In a like manner,  $\beta$ -amino acids such as

may be coupled with 2, the free amine of 8, or 10 to give the corresponding amides which may be converted to the  $\beta$ -amino acid derivatives of compound Ia or Ib following the same chemistry.

Unless otherwise indicated, the term "lower alkyl", "alkyl" or "alk" as employed herein alone or as part of another group includes both straight and branched chain hydrocarbons, containing 1 to 20 carbons, preferably 1 to 10 carbons, more preferably 1 to 8 carbons, in the normal chain, such as methyl, ethyl, propyl, isopropyl, butyl, t-butyl, isobutyl, pentyl, hexyl, isohexyl, heptyl, 4,4-dimethylpentyl, octyl, 2,2,4-trimethyl-pentyl, nonyl, decyl, undecyl, dodecyl,

the various branched chain isomers thereof, and the like as well as such groups including 1 to 4 substituents such as halo, for example F, Br, Cl or I or CF<sub>3</sub>, alkyl, alkoxy, aryl, aryloxy, aryl(aryl) or diaryl, arylalkyl, 5 arylalkyloxy, alkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylalkyloxy, amino, hydroxy, hydroxyalkyl, acyl, heteroaryl, heteroaryloxy, heteroarylalkyl, heteroarylalkoxy, aryloxyalkyl, alkylthio, arylalkylthio, aryloxyaryl, alkylamido, alkanoylamino, 10 arylcarbonylamino, nitro, cyano, thiol, haloalkyl, trihaloalkyl and/or alkylthio.


Unless otherwise indicated, the term "cycloalkyl" as employed herein alone or as part of another group includes saturated or partially unsaturated (containing 1 15 or 2 double bonds) cyclic hydrocarbon groups containing 1 to 3 rings, including monocyclic alkyl, bicyclic alkyl (or bicycloalkyl) and tricyclic alkyl (tricycloalkyl), containing a total of 3 to 20 carbons forming the ring, preferably 3 to 10 carbons, forming the ring and which 20 may be fused to 1 or 2 aromatic rings as described for aryl, which includes cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclodecyl and cyclododecyl, cyclohexenyl, adamantyl,



any of which groups may be optionally substituted with 1 to 4 substituents such as halogen, alkyl, alkoxy, 30 hydroxy, aryl, aryloxy, arylalkyl, cycloalkyl, hydroxyalkyl, alkylamido, alkanoylamino, oxo, acyl, arylcarbonylamino, amino, nitro, cyano, thiol and/or alkylthio and/or any of the substituents for alkyl.

The term "cycloalkenyl" as employed herein alone or as part of another group refers to cyclic hydrocarbons containing 3 to 12 carbons, preferably 5 to 10 carbons and 1 or 2 double bonds. Exemplary cycloalkenyl groups include cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclooctenyl, cyclohexadienyl, and cycloheptadienyl, which may be optionally substituted as defined for cycloalkyl.

The term "cycloalkylene" as employed herein refers to a "cycloalkyl" group which includes free bonds and thus is a linking group such as



and the like, and may optionally be substituted as defined above for "cycloalkyl".

The term "alkanoyl" as used herein alone or as part of another group refers to alkyl linked to a carbonyl group.

Unless otherwise indicated, the term "lower alkenyl" or "alkenyl" as used herein by itself or as part of another group refers to straight or branched chain radicals of 2 to 20 carbons, preferably 2 to 12 carbons, and more preferably 1 to 8 carbons in the normal chain, which include one to six double bonds in the normal chain, such as vinyl, 2-propenyl, 3-butenyl, 2-butenyl, 4-pentenyl, 3-pentenyl, 2-hexenyl, 3-hexenyl, 2-heptenyl, 3-heptenyl, 4-heptenyl, 3-octenyl, 3-nonenyl, 4-decenyl, 3-undecenyl, 4-dodecenyl, 4,8,12-tetradecatrienyl, and the like, and which may be optionally substituted with 1 to 4 substituents, namely, halogen, haloalkyl, alkyl, alkoxy, alkenyl, alkynyl, aryl, arylalkyl, cycloalkyl, amino, hydroxy, heteroaryl, cycloheteroalkyl, alkanoylamino, alkylamido, arylcarbonyl-amino, nitro, cyano, thiol, alkylthio and/or any of the alkyl substituents set out herein.

Unless otherwise indicated, the term "lower alkynyl" or "alkynyl" as used herein by itself or as part of another group refers to straight or branched chain

radicals of 2 to 20 carbons, preferably 2 to 12 carbons and more preferably 2 to 8 carbons in the normal chain, which include one triple bond in the normal chain, such as 2-propynyl, 3-butynyl, 2-butynyl, 4-pentynyl, 3-pentynyl, 2-hexynyl, 3-hexynyl, 2-heptynyl, 3-heptynyl, 4-heptynyl, 3-octynyl, 3-nonynyl, 4-decynyl, 3-undecynyl, 4-dodecynyl and the like, and which may be optionally substituted with 1 to 4 substituents, namely, halogen, haloalkyl, alkyl, alkoxy, alkenyl, alkynyl, aryl, arylalkyl, cycloalkyl, amino, heteroaryl, cycloheteroalkyl, hydroxy, alkanoylamino, alkylamido, arylcarbonylamino, nitro, cyano, thiol, and/or alkylthio, and/or any of the alkyl substituents set out herein.

The terms "arylalkenyl" and "arylalkynyl" as used alone or as part of another group refer to alkenyl and alkynyl groups as described above having an aryl substituent.

Where alkyl groups as defined above have single bonds for attachment to other groups at two different carbon atoms, they are termed "alkylene" groups and may optionally be substituted as defined above for "alkyl".

Where alkenyl groups as defined above and alkynyl groups as defined above, respectively, have single bonds for attachment at two different carbon atoms, they are termed "alkenylene groups" and "alkynylene groups", respectively, and may optionally be substituted as defined above for "alkenyl" and "alkynyl".


The term "halogen" or "halo" as used herein alone or as part of another group refers to chlorine, bromine, fluorine, and iodine as well as  $\text{CF}_3$ , with chlorine or fluorine being preferred.

The term "metal ion" refers to alkali metal ions such as sodium, potassium or lithium and alkaline earth metal ions such as magnesium and calcium, as well as zinc and aluminum.



Unless otherwise indicated, the term "aryl" as employed herein alone or as part of another group refers to monocyclic and bicyclic aromatic groups containing 6 to 10 carbons in the ring portion (such as phenyl or naphthyl including 1-naphthyl and 2-naphthyl) and may optionally include one to three additional rings fused to a carbocyclic ring or a heterocyclic ring (such as aryl, cycloalkyl, heteroaryl or cycloheteroalkyl rings for example

10



and may be optionally substituted through available carbon atoms with 1, 2, or 3 groups selected from hydrogen, halo, haloalkyl, alkyl, haloalkyl, alkoxy, haloalkoxy, alkenyl, trifluoromethyl, trifluoromethoxy, alkynyl, cycloalkylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl, aryl, heteroaryl, arylalkyl, aryloxy, aryloxyalkyl, arylalkoxy, arylthio, arylazo, heteroarylalkyl, heteroarylalkenyl, heteroarylheteroaryl, heteroaryloxy, hydroxy, nitro, cyano, amino, substituted amino wherein the amino includes 1 or 2 substituents (which are alkyl, aryl or any of the other aryl compounds mentioned in the definitions), thiol, alkylthio, arylthio, heteroarylthio, arylthioalkyl, alkoxyarylthio, alkylcarbonyl, arylcarbonyl, alkylaminocarbonyl, arylaminocarbonyl, alkoxy carbonyl, aminocarbonyl, alkylcarbonyloxy, arylcarbonyloxy, alkylcarbonylamino, arylcarbonylamino, arylsulfinyl, arylsulfinylalkyl,

arylsulfonylamino or arylsulfon-aminocarbonyl and/or any of the alkyl substituents set out herein.

Unless otherwise indicated, the term "lower alkoxy", "alkoxy", "aryloxy" or "aralkoxy" as employed  
5 herein alone or as part of another group includes any of the above alkyl, aralkyl or aryl groups linked to an oxygen atom.

Unless otherwise indicated, the term "substituted amino" as employed herein alone or as part of another  
10 group refers to amino substituted with one or two substituents, which may be the same or different, such as alkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl, cycloalkyl, cycloalkylalkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl or  
15 thioalkyl. These substituents may be further substituted with any of the R<sup>1</sup> groups or substituents for R<sup>1</sup> as set out above. In addition, the amino substituents may be taken together with the nitrogen atom to which they are attached to form 1-pyrrolidinyl, 1-piperidinyl, 1-  
20 azepinyl, 4-morpholinyl, 4-thiamorpholinyl, 1-piperazinyl, 4-alkyl-1-piperazinyl, 4-arylalkyl-1-piperazinyl, 4-diarylalkyl-1-piperazinyl, 1-pyrrolidinyl, 1-piperidinyl, or 1-azepinyl, optionally substituted with alkyl, alkoxy, alkylthio, halo, trifluoromethyl or  
25 hydroxy.


Unless otherwise indicated, the term "lower alkylthio", "alkylthio", "arylthio" or "aralkylthio" as employed herein alone or as part of another group includes any of the above alkyl, aralkyl or aryl groups  
30 linked to a sulfur atom.

Unless otherwise indicated, the term "lower alkylamino", "alkylamino", "arylamino", or "arylalkylamino" as employed herein alone or as part of another group includes any of the above alkyl, aryl or  
35 arylalkyl groups linked to a nitrogen atom.

Unless otherwise indicated, the term "acyl" as employed herein by itself or part of another group, as

defined herein, refers to an organic radical linked to a carbonyl ( $\text{C}=\text{O}$ ) group; examples of acyl groups include any of the  $\text{R}^1$  groups attached to a carbonyl, such as alkanoyl, alkenoyl, aroyl, aralkanoyl, heteroaroyl, cycloalkanoyl, cycloheteroalkanoyl and the like.


Unless otherwise indicated, the term "cycloheteroalkyl" as used herein alone or as part of another group refers to a 5-, 6- or 7-membered saturated or partially unsaturated ring which includes 1 to 2 hetero atoms such as nitrogen, oxygen and/or sulfur, linked through a carbon atom or a heteroatom, where possible, optionally via the linker  $(\text{CH}_2)_r$  (where  $r$  is 1, 2 or 3), such as:




and the like. The above groups may include 1 to 4 substituents such as alkyl, halo, oxo and/or any of the alkyl substituents set out herein. In addition, any of the cycloheteroalkyl rings can be fused to a cycloalkyl, aryl, heteroaryl or cycloheteroalkyl ring.

Unless otherwise indicated, the term "heteroaryl" as used herein alone or as part of another group refers to a 5- or 6- membered aromatic ring which includes 1, 2,

- 3 or 4 hetero atoms such as nitrogen, oxygen or sulfur, and such rings fused to an aryl, cycloalkyl, heteroaryl or cycloheteroalkyl ring (e.g. benzothiophenyl, indolyl), and includes possible N-oxides. The heteroaryl group may optionally include 1 to 4 substituents such as any of the substituents set out above for alkyl. Examples of heteroaryl groups include the following:



10



15 and the like.

The term "cycloheteroalkylalkyl" as used herein alone or as part of another group refers to cycloheteroalkyl groups as defined above linked through a C atom or heteroatom to a  $(CH_2)_r$  chain.

20 The term "heteroarylalkyl" or "heteroarylalkenyl" as used herein alone or as part of another group refers to a heteroaryl group as defined above linked through a C atom or heteroatom to a  $-(CH_2)_r-$  chain, alkylene or alkenylene as defined above.

25 The term "polyhaloalkyl" as used herein refers to an "alkyl" group as defined above which includes from 2 to 9, preferably from 2 to 5, halo substituents, such as F or Cl, preferably F, such as  $CF_3CH_2$ ,  $CF_3$  or  $CF_3CF_2CH_2$ .

The term "polyhaloalkoxy" as used herein refers to an "alkoxy" or "alkyloxy" group as defined above which includes from 2 to 9, preferably from 2 to 5, halo substituents, such as F or Cl, preferably F, such as  
5  $\text{CF}_3\text{CH}_2\text{O}$ ,  $\text{CF}_3\text{O}$  or  $\text{CF}_3\text{CF}_2\text{CH}_2\text{O}$ .

All stereoisomers of the compounds of the instant invention are contemplated, either in admixture or in pure or substantially pure form. The compounds of the present invention can have asymmetric centers at any of  
10 the carbon atoms including any one or the R substituents. Consequently, compounds of formula I can exist in enantiomeric or diastereomeric forms or in mixtures thereof. The processes for preparation can utilize racemates, enantiomers or diastereomers as starting  
15 materials. When diastereomeric or enantiomeric products are prepared, they can be separated by conventional methods for example, chromatographic or fractional crystallization.

Where desired, the compounds of structure I may be  
20 used in combination with one or more other types of antidiabetic agents (employed to treat diabetes and related diseases) and/or one or more other types of therapeutic agents which may be administered orally in the same dosage form, in a separate oral dosage form or  
25 by injection.

The other type of antidiabetic agent which may be optionally employed in combination with the DP4 inhibitor of formula I may be 1,2,3 or more antidiabetic agents or antihyperglycemic agents including insulin secretagogues  
30 or insulin sensitizers, or other antidiabetic agents preferably having a mechanism of action different from DP4 inhibition and may include biguanides, sulfonyl ureas, glucosidase inhibitors, PPAR  $\gamma$  agonists, such as thiazolidinediones, SGLT2 inhibitors, PPAR  $\alpha/\gamma$  dual  
35 agonists,  $\alpha_2$  inhibitors, glycogen phosphorylase inhibitors, advanced glycosylation end (AGE) products inhibitors, and/or meglitinides, as well as insulin,

and/or glucagon-like peptide-1 (GLP-1) or mimetics thereof.

It is believed that the use of the compounds of structure I in combination with 1, 2, 3 or more other antidiabetic agents produces antihyperglycemic results greater than that possible from each of these medicaments alone and greater than the combined additive antihyperglycemic effects produced by these medicaments.

The other antidiabetic agent may be an oral antihyperglycemic agent preferably a biguanide such as metformin or phenformin or salts thereof, preferably metformin HCl.

Where the other antidiabetic agent is a biguanide, the compounds of structure I will be employed in a weight ratio to biguanide within the range from about 0.01:1 to about 100:1, preferably from about 0.1:1 to about 5:1.

The other antidiabetic agent may also preferably be a sulfonyl urea such as glyburide (also known as glibenclamide), glimepiride (disclosed in U.S. Patent No. 4,379,785), glipizide, gliclazide or chlorpropamide, other known sulfonylureas or other antihyperglycemic agents which act on the ATP-dependent channel of the  $\beta$ -cells, with glyburide and glipizide being preferred, which may be administered in the same or in separate oral dosage forms.

The compounds of structure I will be employed in a weight ratio to the sulfonyl urea in the range from about 0.01:1 to about 100:1, preferably from about 0.05:1 to about 5:1.

The oral antidiabetic agent may also be a glucosidase inhibitor such as acarbose (disclosed in U.S. Patent No. 4,904,769) or miglitol (disclosed in U.S. Patent No. 4,639,436), which may be administered in the same or in a separate oral dosage forms.

The compounds of structure I will be employed in a weight ratio to the glucosidase inhibitor within the

range from about 0.01:1 to about 100:1, preferably from about 0.2:1 to about 50:1.

The compounds of structure I may be employed in combination with a PPAR  $\gamma$  agonist such as a  
5 thiazolidinedione oral anti-diabetic agent or other insulin sensitizers (which has an insulin sensitivity effect in NIDDM patients) such as troglitazone (Warner-Lambert's Rezulin<sup>®</sup>, disclosed in U.S. Patent No. 4,572,912), rosiglitazone (SKB), pioglitazone (Takeda),  
10 Mitsubishi's MCC-555 (disclosed in U.S. Patent No. 5,594,016), Glaxo-Wellcome's GL-262570, englitazone (CP-68722, Pfizer) or darglitazone (CP-86325, Pfizer, isaglitazone (MIT/J&J), JTT-501 (JPNT/P&U), L-895645 (Merck), R-119702 (Sankyo/WL), NN-2344 (Dr. Reddy/NN), or  
15 YM-440 (Yamanouchi), preferably rosiglitazone and pioglitazone.

The compounds of structure I will be employed in a weight ratio to the thiazolidinedione in an amount within the range from about 0.01:1 to about 100:1, preferably  
20 from about 0.1:1 to about 10:1.

The sulfonyl urea and thiazolidinedione in amounts of less than about 150 mg oral antidiabetic agent may be incorporated in a single tablet with the compounds of structure I.

25 The compounds of structure I may also be employed in combination with a antihyperglycemic agent such as insulin or with glucagon-like peptide-1 (GLP-1) such as GLP-1(1-36) amide, GLP-1(7-36) amide, GLP-1(7-37) (as disclosed in U.S. Patent No. 5,614,492 to Habener,  
30 disclosure of which is incorporated herein by reference), or a GLP-1 mimic such as AC2993 or Exendin-4 (Amylin) and LY-315902 or LY-307167 (Lilly) and NN2211 (Novo-Nordisk), which may be administered via injection, intranasal, or by transdermal or buccal devices.

35 Where present, metformin, the sulfonyl ureas, such as glyburide, glimepiride, glipyrider, glipizide, chlorpropamide and gliclazide and the glucosidase

inhibitors acarbose or miglitol or insulin (injectable, pulmonary, buccal, or oral) may be employed in formulations as described above and in amounts and dosing as indicated in the Physician's Desk Reference (PDR).

- 5           Where present, metformin or salt thereof may be employed in amounts within the range from about 500 to about 2000 mg per day which may be administered in single or divided doses one to four times daily.

- 10           Where present, the thiazolidinedione anti-diabetic agent may be employed in amounts within the range from about 0.01 to about 2000 mg/day which may be administered in single or divided doses one to four times per day.

- 15           Where present insulin may be employed in formulations, amounts and dosing as indicated by the Physician's Desk Reference.

- 20           Where present GLP-1 peptides may be administered in oral buccal formulations, by nasal administration (for example inhalation spray) or parenterally as described in U.S. Patent Nos. 5,346,701 (TheraTech), 5,614,492 and 5,631,224 which are incorporated herein by reference.

- 25           The other antidiabetic agent may also be a PPAR  $\alpha/\gamma$  dual agonist such as AR-HO39242 (Astra/Zeneca), GW-409544 (Glaxo-Wellcome), KRP297 (Kyorin Merck) as well as those disclosed by Murakami et al, "A Novel Insulin Sensitizer Acts As a Coligand for Peroxisome Proliferation - Activated Receptor Alpha (PPAR alpha) and PPAR gamma. Effect on PPAR alpha Activation on Abnormal Lipid Metabolism in Liver of Zucker Fatty Rats", Diabetes 47, 1841-1847 (1998), and in U.S. application Serial No. 09/664,598, filed September 18, 2000, (attorney file LA29NP) the disclosure of which is incorporated herein by reference, employing dosages as set out therein, which compounds designated as preferred are preferred for use herein.

- 35           The other antidiabetic agent may be an SGLT2 inhibitor such as disclosed in U.S. application Serial No. 09/679,027, filed October 4, 2000 (attorney file



LA49NP), which is incorporated herein by reference, employing dosages as set out herein. Preferred are the compounds designated as preferred in the above application.

5           The other antidiabetic agent which may be optionally employed in combination with the DP4 inhibitor of formula I may be an  $\alpha$ P2 inhibitor such as disclosed in U.S. application Serial No. 09/391,053, filed September 7, 1999, and U.S. application Serial No. 09/519,079,  
10       filed March 6, 2000 (attorney file LA27NP), which is incorporated herein by reference, employing dosages as set out herein. Preferred are the compounds designated as preferred in the above application.

          The other antidiabetic agent which may be  
15       optionally employed in combination with the DP4 inhibitor of formula I may be a glycogen phosphorylase inhibitor such as disclosed in WO 96/39384, WO 96/39385, EP 978279, WO 2000/47206, WO 99/43663, and U.S. Patent Nos. 5,952,322 and 5,998,463, WO 99/26659 and EP 1041068.

20       The meglitinide which may optionally be employed in combination with the compound of formula I of the invention may be repaglinide, nateglinide (Novartis) or KAD1229 (PF/Kissei), with repaglinide being preferred.

          The DP4 inhibitor of formula I will be employed in  
25       a weight ratio to the meglitinide, PPAR  $\gamma$  agonist, PPAR  $\alpha/\gamma$  dual agonist, SGLT2 inhibitor,  $\alpha$ P2 inhibitor, or glycogen phosphorylase inhibitor within the range from about 0.01:1 to about 100:1, preferably from about 0.1:1 to about 10:1.

30       The hypolipidemic agent or lipid-modulating agent which may be optionally employed in combination with the compounds of formula I of the invention may include 1,2,3 or more MTP inhibitors, HMG CoA reductase inhibitors, squalene synthetase inhibitors, fibric acid derivatives,  
35       ACAT inhibitors, lipoxxygenase inhibitors, cholesterol absorption inhibitors, ileal  $\text{Na}^+$ /bile acid cotransporter inhibitors, upregulators of LDL receptor activity, ATP


citrate lyase inhibitors, cholesteryl ester transfer protein inhibitors, bile acid sequestrants, and/or nicotinic acid and derivatives thereof.

MTP inhibitors employed herein include MTP inhibitors disclosed in U.S. Patent No. 5,595,872, U.S. Patent No. 5,739,135, U.S. Patent No. 5,712,279, U.S. Patent No. 5,760,246, U.S. Patent No. 5,827,875, U.S. Patent No. 5,885,983 and U.S. Application Serial No. 09/175,180 filed October 20, 1998, now U.S. Patent No. 5,962,440. Preferred are each of the preferred MTP inhibitors disclosed in each of the above patents and applications.

All of the above U.S. Patents and applications are incorporated herein by reference.

Most preferred MTP inhibitors to be employed in accordance with the present invention include preferred MTP inhibitors as set out in U.S. Patent Nos. 5,739,135 and 5,712,279, and U.S. Patent No. 5,760,246 as well as implitapide (Bayer).

The most preferred MTP inhibitor is 9-[4-[4-[[2-(2,2,2-Trifluoroethoxy)benzoyl]amino]-1-piperidinyl]butyl]-N-(2,2,2-trifluoroethyl)-9H-fluorene-9-carboxamide



The hypolipidemic agent may be an HMG CoA reductase inhibitor which includes, but is not limited to, mevastatin and related compounds as disclosed in U.S. Patent No. 3,983,140, lovastatin (mevinolin) and related compounds as disclosed in U.S. Patent No. 4,231,938, pravastatin and related compounds such as disclosed in

U.S. Patent No. 4,346,227, simvastatin and related compounds as disclosed in U.S. Patent Nos. 4,448,784 and 4,450,171. Other HMG CoA reductase inhibitors which may be employed herein include, but are not limited to, 5 fluvastatin, disclosed in U.S. Patent No. 5,354,772, cerivastatin disclosed in U.S. Patent Nos. 5,006,530 and 5,177,080, atorvastatin disclosed in U.S. Patent Nos. 4,681,893, 5,273,995, 5,385,929 and 5,686,104, atavastatin (Nissan/Sankyo's nisvastatin (NK-104)) 10 disclosed in U.S. Patent No. 5,011,930, Shionogi-Astra/Zeneca visastatin (ZD-4522) disclosed in U.S. Patent No. 5,260,440.

The squalene synthetase inhibitors suitable for use herein include, but are not limited to,  $\alpha$ -phosphono- 15 sulfonates disclosed in U.S. Patent No. 5,712,396, those disclosed by Biller et al, J. Med. Chem., 1988, Vol. 31, No. 10, pp 1869-1871, including isoprenoid (phosphinylmethyl)phosphonates as well as other known squalene synthetase inhibitors, for example, as disclosed in U.S. 20 Patent No. 4,871,721 and 4,924,024 and in Biller, S.A., Neuenschwander, K., Ponpipom, M.M., and Poulter, C.D., Current Pharmaceutical Design, 2, 1-40 (1996).

In addition, other squalene synthetase inhibitors suitable for use herein include the terpenoid 25 pyrophosphates disclosed by P. Ortiz de Montellano et al, J. Med. Chem., 1977, 20, 243-249, the farnesyl diphosphate analog A and presqualene pyrophosphate (PSQ-PP) analogs as disclosed by Corey and Volante, J. Am. Chem. Soc., 1976, 98, 1291-1293, phosphinylphosphonates 30 reported by McClard, R.W. et al, J.A.C.S., 1987, 109, 5544 and cyclopropanes reported by Capson, T.L., PhD dissertation, June, 1987, Dept. Med. Chem. U of Utah, Abstract, Table of Contents, pp 16, 17, 40-43, 48-51, Summary.

35 Other hypolipidemic agents suitable for use herein include, but are not limited to, fibric acid derivatives, such as fenofibrate, gemfibrozil, clofibrate,

bezafibrate, ciprofibrate, clinofibrate and the like, probucol, and related compounds as disclosed in U.S. Patent No. 3,674,836, probucol and gemfibrozil being preferred, bile acid sequestrants such as cholestyramine, 5 colestipol and DEAE-Sephadex (Secholex®, Policexide®), as well as lipostabil (Rhone-Poulenc), Eisai E-5050 (an N-substituted ethanolamine derivative), imanixil (HOE-402), tetrahydrolipstatin (THL), istigmastanylphosphorylcholine (SPC, Roche), aminocyclodextrin (Tanabe 10 Seiyoku), Ajinomoto AJ-814 (azulene derivative), melinamide (Sumitomo), Sandoz 58-035, American Cyanamid CL-277,082 and CL-283,546 (disubstituted urea derivatives), nicotinic acid, acipimox, acifran, neomycin, p-aminosalicylic acid, aspirin, 15 poly(diallylmethylamine) derivatives such as disclosed in U.S. Patent No. 4,759,923, quaternary amine poly(diallyldimethylammonium chloride) and ionenes such as disclosed in U.S. Patent No. 4,027,009, and other known serum cholesterol lowering agents.

20           The other hypolipidemic agent may be an ACAT inhibitor such as disclosed in, Drugs of the Future 24, 9-15 (1999), (Avasimibe); "The ACAT inhibitor, Cl-1011 is effective in the prevention and regression of aortic fatty streak area in hamsters", Nicolosi et al, 25 Atherosclerosis (Shannon, Irel). (1998), 137(1), 77-85; "The pharmacological profile of FCE 27677: a novel ACAT inhibitor with potent hypolipidemic activity mediated by selective suppression of the hepatic secretion of ApoB100-containing lipoprotein", Ghiselli, Giancarlo, 30 Cardiovasc. Drug Rev. (1998), 16(1), 16-30; "RP 73163: a bioavailable alkylsulfanyl-diphenylimidazole ACAT inhibitor", Smith, C., et al, Bioorg. Med. Chem. Lett. (1996), 6(1), 47-50; "ACAT inhibitors: physiologic mechanisms for hypolipidemic and anti-atherosclerotic 35 activities in experimental animals", Krause et al, Editor(s): Ruffolo, Robert R., Jr.; Hollinger, Manfred A., Inflammation: Mediators Pathways (1995), 173-98,

Publisher: CRC, Boca Raton, Fla.; "ACAT inhibitors: potential anti-atherosclerotic agents", Sliskovic et al, Curr. Med. Chem. (1994), 1(3), 204-25; "Inhibitors of acyl-CoA:cholesterol O-acyl transferase (ACAT) as hypocholesterolemic agents. 6. The first water-soluble ACAT inhibitor with lipid-regulating activity. Inhibitors of acyl-CoA:cholesterol acyltransferase (ACAT). 7. Development of a series of substituted N-phenyl-N'-[(1-phenylcyclopentyl)methyl]ureas with enhanced hypocholesterolemic activity", Stout et al, Chemtracts: Org. Chem. (1995), 8(6), 359-62, or TS-962 (Taisho Pharmaceutical Co. Ltd).

The hypolipidemic agent may be an upregulator of LD2 receptor activity such as MD-700 (Taisho Pharmaceutical Co. Ltd) and LY295427 (Eli Lilly).

The hypolipidemic agent may be a cholesterol absorption inhibitor preferably Schering-Plough's SCH48461 as well as those disclosed in Atherosclerosis 115, 45-63 (1995) and J. Med. Chem. 41, 973 (1998).

The hypolipidemic agent may be an ileal Na<sup>+</sup>/bile acid cotransporter inhibitor such as disclosed in Drugs of the Future, 24, 425-430 (1999).

The lipid-modulating agent may be a cholesteryl ester transfer protein (CETP) inhibitor such as Pfizer's CP 529,414 (WO/0038722 and EP 818448) and Pharmacia's SC-744 and SC-795.

The ATP citrate lyase inhibitor which may be employed in the combination of the invention may include, for example, those disclosed in U.S. Patent No. 5,447,954.

Preferred hypolipidemic agents are pravastatin, lovastatin, simvastatin, atorvastatin, fluvastatin, cerivastatin, atavastatin and ZD-4522.

The above-mentioned U.S. patents are incorporated herein by reference. The amounts and dosages employed will be as indicated in the Physician's Desk Reference and/or in the patents set out above.

The compounds of formula I of the invention will be employed in a weight ratio to the hypolipidemic agent (were present), within the range from about 500:1 to about 1:500, preferably from about 100:1 to about 1:100.

5       The dose administered must be carefully adjusted according to age, weight and condition of the patient, as well as the route of administration, dosage form and regimen and the desired result.

10       The dosages and formulations for the hypolipidemic agent will be as disclosed in the various patents and applications discussed above.

15       The dosages and formulations for the other hypolipidemic agent to be employed, where applicable, will be as set out in the latest edition of the Physicians' Desk Reference.

20       For oral administration, a satisfactory result may be obtained employing the MTP inhibitor in an amount within the range of from about 0.01 mg/kg to about 500 mg and preferably from about 0.1 mg to about 100 mg, one to four times daily.

25       A preferred oral dosage form, such as tablets or capsules, will contain the MTP inhibitor in an amount of from about 1 to about 500 mg, preferably from about 2 to about 400 mg, and more preferably from about 5 to about 250 mg, one to four times daily.

30       For oral administration, a satisfactory result may be obtained employing an HMG CoA reductase inhibitor, for example, pravastatin, lovastatin, simvastatin, atorvastatin, fluvastatin or cerivastatin in dosages employed as indicated in the Physician's Desk Reference, such as in an amount within the range of from about 1 to 2000 mg, and preferably from about 4 to about 200 mg.

35       The squalene synthetase inhibitor may be employed in dosages in an amount within the range of from about 10 mg to about 2000 mg and preferably from about 25 mg to about 200 mg.

A preferred oral dosage form, such as tablets or capsules, will contain the HMG CoA reductase inhibitor in an amount from about 0.1 to about 100 mg, preferably from about 5 to about 80 mg, and more preferably from about 10 to about 40 mg.

A preferred oral dosage form, such as tablets or capsules will contain the squalene synthetase inhibitor in an amount of from about 10 to about 500 mg, preferably from about 25 to about 200 mg.

The other hypolipidemic agent may also be a lipooxygenase inhibitor including a 15-lipooxygenase (15-LO) inhibitor such as benzimidazole derivatives as disclosed in WO 97/12615, 15-LO inhibitors as disclosed in WO 97/12613, isothiazolones as disclosed in WO 96/38144, and 15-LO inhibitors as disclosed by Sendobry et al "Attenuation of diet-induced atherosclerosis in rabbits with a highly selective 15-lipooxygenase inhibitor lacking significant antioxidant properties", Brit. J. Pharmacology (1997) 120, 1199-1206, and Cornicelli et al, "15-Lipooxygenase and its Inhibition: A Novel Therapeutic Target for Vascular Disease", Current Pharmaceutical Design, 1999, 5, 11-20.

The compounds of formula I and the hypolipidemic agent may be employed together in the same oral dosage form or in separate oral dosage forms taken at the same time.

The compositions described above may be administered in the dosage forms as described above in single or divided doses of one to four times daily. It may be advisable to start a patient on a low dose combination and work up gradually to a high dose combination.

The preferred hypolipidemic agent is pravastatin, simvastatin, lovastatin, atorvastatin, fluvastatin or cerivastatin.

The other type of therapeutic agent which may be optionally employed with the DP4 inhibitor of formula I

may be 1, 2, 3 or more of an anti-obesity agent including a beta 3 adrenergic agonist, a lipase inhibitor, a serotonin (and dopamine) reuptake inhibitor, a thyroid receptor beta drug, an anorectic agent and/or a fatty acid oxidation upregulator.

The beta 3 adrenergic agonist which may be optionally employed in combination with a compound of formula I may be AJ9677 (Takeda/Dainippon), L750355 (Merck), or CP331648 (Pfizer) or other known beta 3 agonists as disclosed in U.S. Patent Nos. 5,541,204, 5,770,615, 5,491,134, 5,776,983 and 5,488,064, with AJ9677, L750,355 and CP331648 being preferred.

The lipase inhibitor which may be optionally employed in combination with a compound of formula I may be orlistat or ATL-962 (Alizyme), with orlistat being preferred.

The serotonin (and dopamine) reuptake inhibitor which may be optionally employed in combination with a compound of formula I may be sibutramine, topiramate (Johnson & Johnson) or axokine (Regeneron), with sibutramine and topiramate being preferred.

The thyroid receptor beta compound which may be optionally employed in combination with a compound of formula I may be a thyroid receptor ligand as disclosed in WO97/21993 (U. Cal SF), WO99/00353 (KaroBio) and GB98/284425 (KaroBio), with compounds of the KaroBio applications being preferred.

The anorectic agent which may be optionally employed in combination with a compound of formula I may be dexamphetamine, phentermine, phenylpropanolamine or mazindol, with dexamphetamine being preferred.

The fatty acid oxidation upregulator which may be optionally employed in combination with the compound of formula I can be famoxin (Genset).

The various anti-obesity agents described above may be employed in the same dosage form with the compound of



formula I or in different dosage forms, in dosages and regimens as generally known in the art or in the PDR.

The infertility agent which may be optionally employed in combination with the DP4 inhibitor of the invention may be 1, 2, or more of clomiphene citrate (Clomid®, Aventis), bromocriptine mesylate (Parlodel®, Novartis), LHRH analogs, Lupron (TAP Pharm.), danazol, Danocrine (Sanofi), progestogens or glucocorticoids, which may be employed in amounts specified in the PDR.

10       The agent for polycystic ovary syndrome which may be optionally employed in combination with the DP4 inhibitor of the invention may be 1, 2, or more of gonadotropin releasing hormone (GnRH), leuprolide (Lupron®), Clomid®, Parlodel®, oral contraceptives or

15       insulin sensitizers such as PPAR agonists, or other conventional agents for such use which may be employed in amounts specified in the PDR.

The agent for treating growth disorders and/or frailty which may be optionally employed in combination with the DP4 inhibitor of the invention may be 1, 2, or more of a growth hormone or growth hormone secretagogue such as MK-677 (Merck), CP-424,391 (Pfizer), and compounds disclosed in U.S. Serial No. 09/506,749 filed February 18, 2000 (attorney docket LA26), as well as

20       selective androgen receptor modulators (SARMs), which is incorporated herein by reference, which may be employed in amounts specified in the PDR, where applicable.

The agent for treating arthritis which may be optionally employed in combination with the DP4 inhibitor of the invention may be 1, 2, or more of aspirin, indomethacin, ibuprofen, diclofenac sodium, naproxen, nabumetone (Relafen®, SmithKline Beecham), tolmetin sodium (Tolectin®, Ortho-McNeil), piroxicam (Feldene®, Pfizer), ketorolac tromethamine (Toradol®, Roche),

30       celecoxib (Celebrex®, Searle), rofecoxib (Vioxx®, Merck) and the like, which may be employed in amounts specified in the PDR.

Conventional agents for preventing allograft rejection in transplantation such as cyclosporin, Sandimmune (Novartis), azathioprine, Immuran (Faro) or methotrexate may be optionally employed in combination  
5 with the DP4 inhibitor of the invention, which may be employed in amounts specified in the PDR.

Conventional agents for treating autoimmune diseases such as multiple sclerosis and immunomodulatory diseases such as lupus erythematosus, psoriasis, for  
10 example, azathioprine, Immuran, cyclophosphamide, NSAIDS such as ibuprofen, cox 2 inhibitors such as Vioxx and Celebrex, glucocorticoids and hydroxychloroquine, may be optionally employed in combination with the DP4 inhibitor of the invention, which may be employed in amounts  
15 specified in the PDR.

The AIDS agent which may be optionally employed in combination with the DP4 inhibitor of the invention may be a non-nucleoside reverse transcriptase inhibitor, a nucleoside reverse transcriptase inhibitor, a protease  
20 inhibitor and/or an AIDS adjunct anti-infective and may be 1, 2, or more of dronabinol (Marinol®, Roxane Labs), didanosine (Videx®, Bristol-Myers Squibb), megestrol acetate (Megace®, Bristol-Myers Squibb), stavudine (Zerit®, Bristol-Myers Squibb), delavirdine mesylate  
25 (Rescriptor®, Pharmacia), lamivudine/zidovudine (Combivir™, Glaxo), lamivudine (Epivir™, Glaxo), zalcitabine (Hivid®, Roche), zidovudine (Retrovir®, Glaxo), indinavir sulfate (Crixivan®, Merck), saquinavir (Fortovase™, Roche), saquinovir mesylate (Invirase®,  
30 Roche), ritonavir (Norvir®, Abbott), nelfinavir (Viracept®, Agouron).

The above anti-AIDS agents may be employed in amounts specified in the PDR.

The agent for treating inflammatory bowel disease  
35 or syndrome which may be optionally employed in combination with the DP4 inhibitor of the invention may be 1, 2, or more of sulfasalazine, salicylates,

mesalamine (Asacol®, P&G) or Zelmac®, (Bristol-Myers Squibb), which may be employed in amounts specified in the PDR or otherwise known in the art.

The agent for treating osteoporosis which may be optionally employed in combination with the DP4 inhibitor of the invention may be 1, 2, or more of alendronate sodium (Fosamax®, Merck, tiludronate (Skelid®, Sanofi), etidronate disodium (Didronel®, P&G), raloxifene HCl (Evista®, Lilly), which may be employed in amounts specified in the PDR.

In carrying out the method of the invention, a pharmaceutical composition will be employed containing the compounds of structure I, with or without another antidiabetic agent and/or other type therapeutic agent, in association with a pharmaceutical vehicle or diluent. The pharmaceutical composition can be formulated employing conventional solid or liquid vehicles or diluents and pharmaceutical additives of a type appropriate to the mode of desired administration. The compounds can be administered to mammalian species including humans, monkeys, dogs, etc. by an oral route, for example, in the form of tablets, capsules, granules or powders, or they can be administered by a parenteral route in the form of injectable preparations. The dose for adults is preferably between 10 and 1,000 mg per day, which can be administered in a single dose or in the form of individual doses from 1-4 times per day.

A typical capsule for oral administration contains compounds of structure I (250 mg), lactose (75 mg) and magnesium stearate (15 mg). The mixture is passed through a 60 mesh sieve and packed into a No. 1 gelatin capsule.

A typical injectable preparation is produced by aseptically placing 250 mg of compounds of structure I into a vial, aseptically freeze-drying and sealing. For use, the contents of the vial are mixed with 2 mL of

physiological saline, to produce an injectable preparation.

DP4 inhibitor activity of the compounds of the invention may be determined by use of an in vitro assay system which measures the potentiation of inhibition of DP4. Inhibition constants ( $K_i$  values) for the DP4 inhibitors of the invention may be determined by the method described below.

#### 10 Purification of Porcine Dipeptidyl Peptidase IV

Porcine enzyme was purified as previously described (1), with several modifications. Kidneys from 15-20 animals were obtained, and the cortex was dissected away and frozen at  $-80^{\circ}\text{C}$ . Frozen tissue (2000 -2500 g) was homogenized in 12 L of 0.25 M sucrose in a Waring blender. The homogenate then was left at  $37^{\circ}\text{C}$  for 18 hours to facilitate cleavage of DP-4 from cell membranes. After the cleavage step, the homogenate was clarified by centrifugation at 7000 X g for 20 min at  $4^{\circ}\text{C}$ , and the supernatant was collected. Solid ammonium sulfate was added to 60% saturation, and the precipitate was collected by centrifugation at 10,000 X g and was discarded. Additional ammonium sulfate was added to the supernatant to 80% saturation, and the 80% pellet was collected and dissolved in 20 mM  $\text{Na}_2\text{HPO}_4$ , pH 7.4.

After dialysis against 20 mM  $\text{Na}_2\text{HPO}_4$ , pH 7.4, the preparation was clarified by centrifugation at 10,000 X g. The clarified preparation then was applied to 300 mL of ConA Sepharose that had been equilibrated in the same buffer. After washing with buffer to a constant  $A_{280}$ , the column was eluted with 5% (w/v) methyl  $\alpha$ -D-mannopyranoside. Active fractions were pooled, concentrated, and dialyzed against 5 mM sodium acetate, pH 5.0. Dialyzed material then was flowed through a 100 mL Pharmacia Resource S column equilibrated in the same buffer. The flow through material was collected and contained most of the enzyme activity. Active material

again was concentrated and dialyzed into 20 mM  $\text{Na}_2\text{HPO}_4$ , pH 7.4. Lastly, the concentrated enzyme was chromatographed on a Pharmacia S-200 gel filtration column to removed low molecular weight contaminants. Purity of column fractions was analyzed by reducing SDS-PAGE, and the purest fractions were pooled and concentrated. Purified enzyme was stored in 20% glycerol at  $-80^\circ\text{C}$ .

#### Assay of Porcine Dipeptidyl Peptidase IV

Enzyme was assayed under steady-state conditions as previously described (2) with gly-pro-p-nitroanilide as substrate, with the following modifications. Reactions contained, in a final volume of 100  $\mu\text{l}$ , 100 mM Aces, 52 mM TRIS, 52 mM ethanolamine, 500  $\mu\text{M}$  gly-pro-p-nitroanilide, 0.2 % DMSO, and 4.5 nM enzyme at  $25^\circ\text{C}$ , pH 7.4. For single assays at 10  $\mu\text{M}$  test compound, buffer, compound, and enzyme were added to wells of a 96 well microtiter plate, and were incubated at room temperature for 5 min. Reactions were started by addition of substrate. The continuous production of p-nitroaniline was measured at 405 nM for 15 min using a Molecular Devices Tmax plate reader, with a read every 9 seconds. The linear rate of p-nitroaniline production was obtained over the linear portion of each progress curve. A standard curve for p-nitroaniline absorbance was obtained at the beginning of each experiment, and enzyme catalyzed p-nitroaniline production was quantitated from the standard curve. Compounds giving greater than 50% inhibition were selected for further analysis.

For analysis of positive compounds, steady-state kinetic inhibition constants were determined as a function of both substrate and inhibitor concentration. Substrate saturation curves were obtained at gly-pro-p-nitroanilide concentrations from 60  $\mu\text{M}$  to 3600  $\mu\text{M}$ . Additional saturation curves also were obtained in the presence of inhibitor. Complete inhibition experiments contained 11 substrate and 7 inhibitor concentrations,

with triplicate determinations across plates. For tight binding inhibitors with  $K_i$ s less than 20 nM, the enzyme concentration was reduced to 0.5 nM and reaction times were increased to 120 min. Pooled datasets from the  
5 three plates were fitted to the appropriate equation for either competitive, noncompetitive or uncompetitive inhibition.

(1) Rahfeld, J. Schutkowski, M., Faust, J.,  
10 Neubert., Barth, A., and Heins, J. (1991) Biol. Chem. Hoppe-Seyler, 372, 313-318.


(2) Nagatsu, T., Hino, M., Fuyamada, H., Hayakawa, T., Sakakibara, S., Nakagawa, Y., and Takemoto, T. (1976)  
15 Anal. Biochem., 74, 466-476.

The following abbreviations are employed in the Examples and elsewhere herein:

20 Ph = phenyl  
Bn = benzyl  
i-Bu = iso-butyl  
Me = methyl  
Et = ethyl  
25 Pr = propyl  
Bu = butyl  
TMS = trimethylsilyl  
Fmoc = fluorenylmethoxycarbonyl  
Boc or BOC = tert-butoxycarbonyl  
30 Cbz = carbobenzyloxy or carbobenzoxy or benzyloxycarbonyl  
HOAc or AcOH = acetic acid  
DMF = N,N-dimethylformamide  
EtOAc = ethyl acetate  
THF = tetrahydrofuran  
35 TFA = trifluoroacetic acid  
Et<sub>2</sub>NH = diethylamine  
NMM = N-methyl morpholine


- n*-BuLi = *n*-butyllithium  
Pd/C = palladium on carbon  
PtO<sub>2</sub> = platinum oxide  
TEA = triethylamine
- 5 EDAC = 3-ethyl-3'-(dimethylamino)propyl-carbodiimide  
hydrochloride (or 1-[(3-(dimethylamino)propyl)]-3-  
ethylcarbodiimide hydrochloride)  
HOBT or HOBT•H<sub>2</sub>O = 1-hydroxybenzotriazole hydrate  
HOAT = 1-hydroxy-7-azabenzotriazole
- 10 PyBOP reagent = benzotriazol-1-yloxy-tripyrrolidino  
phosphonium hexafluorophosphate  
min = minute(s)  
h or hr = hour(s)  
L = liter
- 15 mL = milliliter  
μL = microliter  
g = gram(s)  
mg = milligram(s)  
mol = mole(s)
- 20 mmol = millimole(s)  
meq = milliequivalent  
rt = room temperature  
sat or sat'd = saturated  
aq. = aqueous
- 25 TLC = thin layer chromatography  
HPLC = high performance liquid chromatography  
LC/MS = high performance liquid chromatography/mass  
spectrometry  
MS or Mass Spec = mass spectrometry
- 30 NMR = nuclear magnetic resonance  
mp = melting point

The following Examples represent preferred  
35 embodiments of the invention.

Example 1

Step 1.

5




Step 1 title compound was synthesized by following the literature procedure [Stephen Hanessian, Ulrich Reinhold, Michel Saulnier, and Stephen Claridge; Bioorganic & Medicinal Chemistry Letters 8 (1998) 2123-2128] or with the following modifications. L-pyroglutamic acid ethyl ester was N-protected as the t-butylcarbamate (Boc<sub>2</sub>O, DMAP or NaH) and then dehydrated to the 4,5-dehydroproline ethyl ester in one pot by carbonyl reduction (triethylborohydride, toluene, -78°C) followed by dehydration (TFAA, lutidine). The title compound was obtained by cyclopropanation of the 4,5-dehydroproline ethyl ester (Et<sub>2</sub>Zn, ClCH<sub>2</sub>I, 1,2-dichloroethane, -15°C). A more detailed protocol is as follows:

Synthesis of 4,5-dehydro-L-proline ethyl ester: L-pyroglutamic acid ethyl ester (200 g, 1.27 mol) was dissolved in 1.2 liters of methylene chloride and treated sequentially with di-tert-butylidicarbonate (297 g, 1.36 mol) and a catalytic DMAP (1.55 g, 0.013 mol) at ambient temperature. After 6 h, the mixture was quenched with



- saturated brine and the organic phase was dried ( $\text{Na}_2\text{SO}_4$ ) and filtered through a short silica gel column to give 323 g (100%) of N-Boc- L-pyroglutamic acid ethyl ester. N-Boc- L-pyroglutamic acid ethyl ester ( 160 g, 0.62 mol)
- 5 was dissolved in 1 liter of toluene, cooled to  $-78^\circ\text{C}$  and treated with lithium triethylborohydride (666 mL of a 1.0 M soln in THF) and added dropwise over 90 minutes. After 3 h, 2,6-lutidine (423 mL, 3.73 mol) was added dropwise followed by DMAP (0.2 g, 0.0016 mol). To this mixture
- 10 was added TFAA (157 g, 0.74 mol) and the reaction was allowed to come to ambient temperature over 2 h. The mixture was diluted with EtOAc and water and the organics were washed with 3 N HCl, water, aqueous bicarbonate and brine and dried ( $\text{Na}_2\text{SO}_4$ ) and filtered through a silica gel
- 15 plug to give 165 g of the crude 4,5-dehydroproline ethyl ester that was purified by flash column chromatography on silica gel with 1:5 ethyl acetate:hexanes to give 120 g, 75% of the olefin.
- 20 Cyclopropanation of 4,5-dehydro-L-proline ethyl ester:  
4,5-Dehydro-L-proline ethyl ester (35.0 g, 0.145 mol) was added to a solution of neat  $\text{Et}_2\text{Zn}$  (35.8 g, 0.209 mol) in 1 liter of 1,2-dichloroethane at  $-15^\circ\text{C}$ . To this mixture was added a dropwise addition of  $\text{ClCH}_2\text{I}$  (102 g, 0.58 mol) over
- 25 1 h and the mixture stirred at  $-15^\circ\text{C}$  for 18 h. The reaction was quenched with saturated aqueous bicarbonate and the solvent was evaporated and the reaction was taken up in EtOAc, washed with brine and purified by silica gel chromatography using a stepwise gradient of from 20%
- 30 EtOAc/hexanes to 50% EtOAc/hexanes to give 17.5 g (50%) of diastereomerically pure step 1 title compound.


Step 2.



To a stirred solution of Step 1 compound (411 mg, 1.61 mmol) in  $\text{CH}_2\text{Cl}_2$  (1.5 mL) at rt was added TFA (1.5 mL).


- 5 The reaction mixture was stirred at rt for 2 h and evaporated. The residue was diluted with  $\text{CH}_2\text{Cl}_2$  and then evaporated and re-evaporated three times to give the title compound as a colorless oil, 433 mg, 100% yield.

10 Step 3.




- To a stirred solution of (S)-N-tert-butoxycarbonyl-isoleucine (372.6 mg, 1.61 mmol) and benzotriazol-1-yloxytripyrrolidinophosphonium hexafluorophosphate (1.25 g, 2.42 mmol) in  $\text{CH}_2\text{Cl}_2$  (6 mL) under nitrogen at rt was
- 15 added 4-methylmorpholine (NMM) (0.36 mL, 3.2 mmol). After 5 min, a solution of Step 2 compound (433 mg, 1.61 mmol) and NMM (0.27 mL, 2.4 mmol) in  $\text{CH}_2\text{Cl}_2$  (1 mL) was added. After addition, the reaction mixture was stirred
- 20 under nitrogen at room temperature overnight. The reaction mixture was diluted with  $\text{CH}_2\text{Cl}_2$  (40 mL) and washed with 4%  $\text{KHSO}_4$  (10 mL), aqueous  $\text{NaHCO}_3$  (10 mL) and brine (10 mL), dried ( $\text{Na}_2\text{SO}_4$ ) and evaporated.
- Purification by flash chromatography (1:4 EtOAc/hexane)
- 25 gave the title compound as a colorless oil, 530 mg, 89% yield.

## Step 4



To a stirred solution of Step 3 compound (530 mg, 1.44 mmol) in MeOH (4 mL) and H<sub>2</sub>O (4 mL) at rt was added LiOH-H<sub>2</sub>O (91 mg, 2.16 mmol). The reaction mixture was stirred at rt overnight and evaporated. Water (10 mL) was added to the residue and extracted with Et<sub>2</sub>O (2 x 10 mL). The aqueous layer was acidified to ~pH 4 by adding 4% KHSO<sub>4</sub> dropwise. The milky solution was extracted with EtOAc (15 mL x 3). Combined EtOAc layers were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated to give the title compound as a white solid, 440 mg, 90% yield.


## Step 5



To a stirred solution of Step 4 compound (300 mg, 0.88 mmol) in THF (6 mL) at -15°C under nitrogen, was added 4-methylmorpholine (0.12 mL, 1.06 mmol) and then isobutyl chloroformate (0.13 mL, 0.97 mmol) over 2 min. White precipitate was formed. The reaction mixture was stirred at -15°C under nitrogen for 25 min and a solution of NH<sub>3</sub> in dioxane (8.8 mL, 4.4 mmol) was added. The reaction mixture was stirred at -15°C for 30 min, warmed to rt and stirred at rt overnight. The reaction mixture was quenched by 4% KHSO<sub>4</sub> to ~pH 4 and extracted with EtOAc (20 mL x 3). The extracts were combined, washed with brine


(10 mL) dried ( $\text{Na}_2\text{SO}_4$ ) and evaporated. Purification by flash column chromatography (1:1 EtOAc/hexane) gave the title compound as a white foam, 268 mg, 90% yield.

## 5 Step 6



To a stirred solution of Step 5 compound (248 mg, 1.38 mmol) and imidazole (94 mg, 1.38 mmol) in dry pyridine (12 mL) at  $-35^\circ\text{C}$  under nitrogen was added  $\text{POCl}_3$  (0.26 mL, 2.76 mmol) dropwise. The reaction mixture was stirred between  $-35^\circ\text{C}$  to  $-20^\circ\text{C}$  for 1 h and evaporated.  $\text{CH}_2\text{Cl}_2$  (10 mL) was added and white precipitates were formed. After filtration, the filtrate was concentrated and purified by flash chromatography (2:5 EtOAc/hexane) to give the title compound as a colorless oil, 196 mg, 88% yield.

## Step 7




To a stirred solution of Step 6 compound (130 mg, 0.4 mmol) in  $\text{CH}_2\text{Cl}_2$  (2 mL) at rt was added TFA (2 mL). The reaction mixture was stirred at rt for 2 h. The reaction mixture was added slowly to a pre-cooled slurry of  $\text{NaHCO}_3$  (3.8 g) in  $\text{H}_2\text{O}$  (3 mL). The mixture was extracted with  $\text{CH}_2\text{Cl}_2$  (6 mL x 5), and the combined  $\text{CH}_2\text{Cl}_2$  layers were evaporated and purified by preparative HPLC to give the title compound as a white powder, 77 mg, 57% yield, mp =


141-143°C. LC/MS gave the correct molecular ion  $[(M+H)^+ = 222]$  for the desired compound.

Example 2

5




Step 1




10 Step 1 title compound was synthesized by following the literature procedure. [Stephen Hanessian, Ulrich Reinhold, Michel Saulnier, and Stephen Claridge; Bioorganic & Medicinal Chemistry Letters 8 (1998) 2123-2128.]


15 Step 2



The title compound was prepared from Step 1 compound, employing the same procedure as that described for Example 1, Steps 2-6. LC/MS gave the correct molecular  
20 ion  $[(M+H)^+ = 222]$  for the desired compound.

Example 3


## Step 1



5


Step 1 title compound was prepared by following the literature procedure. [Willy D. Kollmeyer, U.S. Patent 4,183,857.]

## 10 Step 2




To a stirred solution of (S)-N-tert-butoxycarbonyl-isoleucine (231 mg, 1 mmol) and benzotriazol-1-yloxytripyrrolidinophosphonium hexafluorophosphate (780 mg, 1.5 mmol) in  $\text{CH}_2\text{Cl}_2$  (6 mL) under nitrogen at rt was added 4-methylmorpholine (0.33 mL, 3 mmol). After 5 min, Step 1 compound (120 mg, 1 mmol) was added in one portion. The reaction mixture was stirred under nitrogen at rt overnight and then diluted with  $\text{CH}_2\text{Cl}_2$  (30 mL), washed with 4.1%  $\text{KHSO}_4$  (10 mL), aqueous  $\text{NaHCO}_3$  (10 mL), brine (10 mL), dried ( $\text{Na}_2\text{SO}_4$ ) and evaporated. Purification by flash chromatography on silica gel (2.4 x 20 cm column, 1:3 EtOAc/hexane) gave the title compound as a colorless oil, 290 mg, 90% yield. LC/MS gave the correct molecular ion  $[(\text{M}+\text{H})^+ = 297]$  for the desired compound.

## Step 3



The reaction mixture of Step 2 compound (220 mg, 0.74 mmol) and 4 M HCl in dioxane (1.5 mL, 6 mmol) was stirred at rt for 2 h and evaporated under reduced pressure. Et<sub>2</sub>O was added to the residue and a precipitate was formed. Et<sub>2</sub>O was decanted and this was done three times. The precipitate was dried in vacuo to give the title compound as a white powder, 130 mg (76% yield), mp 205-206°C. LC/MS gave the correct molecular ion [(M+H)<sup>+</sup> = 197] for the desired compound.

## Examples 4-4A




## Step 1




Step 1 title compound, as a 1:1 ratio of enantiomers, was prepared by following the literature procedure. [Willy D. Kollmeyer, U.S. Patent 4,183,857.]

## Step 2



A slurry of (S)-N-tert-butoxycarbonyl-isoleucine (92.5 mg, 0.4 mmol), 1-[(3-(dimethylamino)propyl)-3-ethylcarbodiimide (77 mg, 0.4 mmol) and HOAT (54.4 mg, 0.4 mmol) in  $\text{ClCH}_2\text{CH}_2\text{Cl}$  (0.3 mL) was stirred under nitrogen at rt for 1 h, then Step 1 compound (22 mg, 0.2 mmol) was added, followed by  $\text{Et}_3\text{N}$  (0.015 mL, 0.1 mmol). The reaction mixture was stirred under nitrogen at rt overnight and then diluted with  $\text{CH}_2\text{Cl}_2$  (3 mL), washed with  $\text{H}_2\text{O}$  (1 mL), aqueous  $\text{NaHCO}_3$  (1 mL) and brine (1 mL), dried ( $\text{Na}_2\text{SO}_4$ ) and evaporated. Purification by flash chromatography on silica gel (2.4 x 12 cm column, 2:7 EtOAc/hexane) gave the title compound as a colorless oil, 33 mg, 51% yield. LC/MS gave the correct molecular ion  $[(M+H)^+ = 322]$  for the desired compound.

## Step 3




To a stirred solution of Step 2 compound (30 mg, 0.4 mmol) in  $\text{CH}_2\text{Cl}_2$  (0.5 mL) at rt was added TFA (0.5 mL). The reaction mixture was stirred at rt for 2 h. The reaction mixture was added slowly to a precooled slurry of  $\text{NaHCO}_3$  (0.8 g) in  $\text{H}_2\text{O}$  (1 mL). The mixture was extracted with  $\text{CH}_2\text{Cl}_2$  (2 mL x 5), and combined  $\text{CH}_2\text{Cl}_2$  layers were evaporated and purified by preparative HPLC to give the title compounds as a 1:1 ratio of



diastereomers, 22 mg, 73% yield. LC/MS gave the correct molecular ion  $[(M+H)^+ = 222]$  for the desired compounds.

### Examples 5-5A

5




Step 1



To a solution of Example 4, Step 1 compound (150 mg, 1.39 mmol) in 2-propanol (0.8 mL), was added NaCN (40 mg, 1.0 mmol). The reaction mixture was heated to reflux for 3 h. After cooling to rt, the reaction mixture was evaporated and then slurried in Et<sub>2</sub>O (5 mL). After filtration, the filtrate was evaporated to give Example 4 Step 1 compounds and Example 5 Step 1 compounds (140 mg, 93%) as a 2:1 mixture of diastereomers, each as a racemic mixture.

Step 2



20


A slurry of (S)-N-tert-butoxycarbonyl-isoleucine (595 mg, 2.57 mmol), 1-[(3-(dimethylamino)propyl]-3-ethylcarbodiimide (493 mg, 2.57 mmol) and 1-hydroxy-7-azabenzotriazole (350 mg, 2.57 mmol) in ClCH<sub>2</sub>CH<sub>2</sub>Cl (2 mL) was stirred under nitrogen at rt for 1 h, then Step 1 compound mixture (139 mg, 1.28 mmol) was added. The

reaction mixture was stirred under nitrogen at rt overnight and then diluted with CH<sub>2</sub>Cl<sub>2</sub> (30 mL), washed with H<sub>2</sub>O (10 mL), saturated aqueous NaHCO<sub>3</sub> (10 mL) and brine (10 mL), dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated.


- 5 Purification by flash chromatography on silica gel (2.4 x 20 cm column, 1:3 EtOAc/hexane) gave the Example 4, Step 2 compound (260 mg), and the title compounds (105 mg) as a ratio of 1:1 diastereomers. LC/MS gave the correct molecular ion [(M+H)<sup>+</sup> = 322] for the desired compounds.

10

## Step 3



and




- To a stirred solution of Step 2 compounds (104 mg, 0.32 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (1 mL) at rt was added TFA (1 mL). The reaction mixture was stirred at rt for 2 h. The reaction mixture was added slowly to a precooled slurry of NaHCO<sub>3</sub> (2 g) in H<sub>2</sub>O (2 mL). The mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub> (4 mL x 4), and combined CH<sub>2</sub>Cl<sub>2</sub> layers were evaporated and purified by preparative HPLC to give the title compound Example 5 (36 mg) and Example 5A (36 mg). LC/MS gave the correct molecular ion [(M+H)<sup>+</sup> = 222] for the desired compounds.

Example 6

- 25 **General Method A:** Parallel array synthesis methods for preparation of inhibitors from commercially available amino acids. As shown in Scheme 3, the ester 11, described in Example 1 Step 1, was saponified to the acid with LiOH in THF/H<sub>2</sub>O and converted to the amide 12 by treatment with isobutyl chloroformate/NMM followed by ammonia in dioxane. The Boc protecting group was removed under acidic conditions using TFA in methylene chloride to give 13. The TFA salt was coupled to Boc-t-


butylglycine using either EDAC/HOBT/DMF or EDAC/DMAP/CH<sub>2</sub>Cl<sub>2</sub> to give 14. The amide was dehydrated to the nitrile 15 using POCl<sub>3</sub>/imidazole in pyridine at -20°C and finally deprotected with TFA in CH<sub>2</sub>Cl<sub>2</sub> at ambient temperature to afford the target 16.

**Scheme 3, General Method A (Examples 6-27)**




a. LiOH in THF/H<sub>2</sub>O or MeOH/H<sub>2</sub>O b. *i*-BuOCOCi/ NMM or *i*-BuOCOCi/TEA at -30°C or EDAC, then NH<sub>3</sub> in dioxane or Et<sub>2</sub>O at RT c. TFA, CH<sub>2</sub>Cl<sub>2</sub>, RT d. Boc-L-butylglycine and PyBop/ NMM or EDAC, DMAP, CH<sub>2</sub>Cl<sub>2</sub> e. POCl<sub>3</sub>, pyridine, imidazole, -20°C f. TFA, CH<sub>2</sub>Cl<sub>2</sub>, RT

10



Step 1




15

To a stirred solution of Example 1 Step 1 compound (1.40 g, 5.49 mmol) in 40 mL of a 1:1 methanol:water solution at rt was added lithium hydroxide (0.20 g, 8.30 mmol). The reaction mixture was stirred at rt for 18 h and then


heated to 50°C for 2 h. The mixture was diluted with equal volumes of ether and water (50 mL) and then acidified with KHSO<sub>4</sub> to pH 3. The milky solution was extracted with ether (3 X 20 mL). The combined ether layers were dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated. The residue was stripped from toluene (2 X 10 mL) and dried under reduced pressure to give the title compound as a thick syrup, 1.20 g, 96%.

## 10 Step 2




To a stirred solution of Step 1 compound (1.20 g, 5.28 mmol) in THF (20 mL) at -15°C under nitrogen was added 4-methylmorpholine (0.71 mL, 6.50 mmol) and then isobutyl chloroformate (0.78 mL, 6.00 mmol) over 5 min. The reaction was stirred at -15°C for 30 min, cooled to -30°C and treated with a solution of NH<sub>3</sub> in dioxane (50 mL, 25 mmol). The reaction mixture was stirred at -30°C for 30 min, warmed to rt and stirred overnight. The reaction mixture was quenched with citric acid solution (pH 4) and extracted with ether (3 X 50 mL). The combined organic fractions were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. Purification by flash column chromatography on silica gel with EtOAc gave the Step 2 compound, 1.00 g, 84%.

## Step 3



To a stirred solution of Step 2 compound (0.90 g, 4.00 mmol) in  $\text{CH}_2\text{Cl}_2$  (3 mL) at  $0^\circ\text{C}$  was added TFA (3 mL). The reaction mixture was stirred at  $0^\circ\text{C}$  for 18 h. The reaction mixture was concentrated under reduced pressure to produce title compound in the form of a thick oil, 0.98 g, 100%. The oil gradually solidified upon prolonged standing.


## Step 4



An oven-dried 15-mL test tube was charged with Step 3 compound (56 mg, 0.22 mmol), N-tert-butoxycarbonyl-(L)-tert-leucine (53 mg, 0.23 mmol), dimethylaminopyridine (0.11 g, 0.88 mmol), and  $\text{CH}_2\text{Cl}_2$  (4 mL). The tube was sealed under nitrogen atmosphere and treated with 1-[(3-(dimethylamino)propyl)-3-ethylcarbodiimide (84 mg, 0.44 mmol). The mixture was placed in a shaker and vortexed overnight. The product was purified by solid phase extraction using a United Technology SCX column (2 g of sorbent in a 6 mL column) by loading the material on a SCX ion exchange column and successively washing with  $\text{CH}_2\text{Cl}_2$  (5 mL), 30% methanol in  $\text{CH}_2\text{Cl}_2$  (5 mL), 50% methanol in  $\text{CH}_2\text{Cl}_2$  (5 mL) and methanol (10 mL). The product containing fractions were concentrated under reduced pressure to give the desired amide. Further purification by reverse phase preparative column chromatography on a YMC S5 ODS 20 X 250 mm column gave the title compound, 50 mg (68% yield). Purification conditions: Gradient elution from 30% methanol/water/0.1 TFA to 90% methanol/water/0.1 TFA over 15 min. 5 min. hold at 90%


methanol/water/0.1 TFA. Flow rate: 20 mL/min. Detection wavelength: 220. Retention Time: 14 min.

## Step 5




- 5 An oven-dried 15-mL test tube was charged with Step 4 compound (50 mg, 0.15 mmol), imidazole (31 mg, 0.46 mmol), and pyridine (1 mL). The tube was sealed under nitrogen atmosphere and cooled to  $-30^{\circ}\text{C}$ . Slow addition of
- 10  $\text{POCl}_3$  (141 mg, 88  $\mu\text{L}$ , 0.92 mmol) gave after mixing a thick slurry. The tube was mixed at  $-30^{\circ}\text{C}$  for 3 h and the volatiles evaporated. The product was purified by solid phase extraction using a United Technology silica
- 15 extraction column (2 g of sorbent in a 6 mL column) by loading the material on a silica column and successively washing with  $\text{CH}_2\text{Cl}_2$  (5 mL), 5% methanol in  $\text{CH}_2\text{Cl}_2$  (5 mL), 7% methanol in  $\text{CH}_2\text{Cl}_2$  (5 mL) and 12% methanol in  $\text{CH}_2\text{Cl}_2$  (10 mL). The product containing fractions were pooled and concentrated under reduced pressure to give the title
- 20 compound, 46 mg, 96%.

## Step 6












- 25 An oven-dried 15-mL test tube was charged with Step 5 compound (0.45 mg, 0.14 mmol),  $\text{CH}_2\text{Cl}_2$  (1 mL), and TFA (1 mL). The reaction mixture was vortexed for 40 min at rt,





- diluted with toluene (4 mL) and concentrated under reduced pressure to a thick oil. The product was purified by reverse phase preparative column chromatography on a YMC S5 ODS 20 X 250 mm column to give
- 5 the Example 6 compound, 14 mg, 35%. Purification conditions: gradient elution from 10% methanol/water/0.1 TFA to 90% methanol/water/0.1 TFA over 18 min; 5 min hold at 90% methanol/water/0.1 TFA. Flow rate: 20 mL/min. Detection wavelength: 220. Retention Time: 10 min.
- 10 Examples 7-27 were prepared from amino acids available from commercial sources according to the procedure in Example 6.


Table 1

Example	R	[M + H]
7	<p>Chemical structure of R for Example 7: A side chain with a benzyl group, a thioether, an amine, and a ketone.</p>	302
8	<p>Chemical structure of R for Example 8: A side chain with an indol-3-ylmethyl group, an amine, and a ketone.</p>	295
9	<p>Chemical structure of R for Example 9: A side chain with a methylthio group, an amine, and a ketone.</p>	240
10	<p>Chemical structure of R for Example 10: A side chain with an isobutyl group, an amine, and a ketone.</p>	222
11	<p>Chemical structure of R for Example 11: A side chain with a butyl group, an amine, and a ketone.</p>	222
12	<p>Chemical structure of R for Example 12: A side chain with an isopropyl group, a secondary amine, and a ketone.</p>	222
13	<p>Chemical structure of R for Example 13: A side chain with an isopropyl group, a primary amine, and a ketone.</p>	208
14	<p>Chemical structure of R for Example 14: A side chain with a benzyl group, an amine, and a ketone.</p>	270




15		222
16		206
17		256
18		268
19		220
20		220
21		210
22		262
23		242

24		210
25		281
26		281
27		272


Example 27

## 5 Step 1



(2S,4S,5S)-4,5-methano-L-proline carboxylamide, TFA salt  
 10 (53 mg, 0.22 mmol) was coupled to N-Boc-L-Tyrosine-benzyl  
 ether (82 mg, 0.22 mmol) using PyBop (172 mg, 0.33 mmol)  
 and N-methylmorpholine (67 mg, 0.66 mmol) in 4 mL CH<sub>2</sub>Cl<sub>2</sub>.  
 The reaction stirred for 16 h, was taken up in EtOAc,  
 washed with H<sub>2</sub>O, 1N aqueous HCl, brine, then evaporated  
 15 and purified by silica gel flash chromatography to give  
 the coupled product (FAB MH<sup>+</sup> 480).


## Step 2



5

The Step 1 amide was dehydrated to the nitrile using the general method C (which follows Example 29) (FAB MH+ 462).


## 10 Step 3



The Step 2 benzyl ether was cleaved by catalytic hydrogenolysis using 10% palladium on carbon and 1 atmosphere hydrogen gas in MeOH at rt for 1.5 h. The reaction was filtered through celite and concentrated to an oil and taken on without further purification (FAB MH+ 372).


20

## Step 4



Step 3 N-[N-Boc-L-Tyrosine-]-(2S,4S,5S)-2-cyano-4,5-methano-L-prolylamide was dissolved in CH<sub>2</sub>Cl<sub>2</sub> and TFA was added at rt. The reaction stirred for 1 h and was evaporated and purified by preparative HPLC as described in general method B (set out following Example 29) to afford the title compound (FAB MH+ 272).

Example 28




10


The title compound was prepared by coupling (2S,4S,5S)-4,5-methano-L-proline carboxylamide, TFA salt described in Example 6 Step 3 compound with N-(tert-butyloxy-carbonyl)hydroxyvaline. After hydroxyl protection with triethylsilyl chloride and dehydration of the amide with POCl<sub>3</sub>/imidazole in pyridine and deprotection (N-terminal nitrogen and valine hydroxyl) with TFA using general method C (FAB MH+ 224), the title compound was obtained.

20

Example 29



Step 1




25

N-Boc-L-homoserine (1.20 g, 5.47 mmol) upon treatment with tert-butyldimethylsilyl chloride (1.67 g, 11.04

mmol) and imidazole (938 mg, 13.8 mmol) in THF (17 mL) was stirred as thick slurry for 48 h under N<sub>2</sub>. The solvent was evaporated, and the crude material was dissolved in MeOH (10 mL). The resulting solution was stirred at rt for 2 h. The solvent was evaporated, and the crude material was diluted with CH<sub>2</sub>Cl<sub>2</sub> (50 mL) and treated with 0.1N HCl (2x10 mL). The CH<sub>2</sub>Cl<sub>2</sub> layer was washed with brine and dried over MgSO<sub>4</sub>. Removal of the volatiles gave title compound as an oil (1.8 g), which was used without further purification (LC/Mass, + ion): 334 (M+H).


## Step 2



15


To a stirred solution of Step 1 compound (333 mg, 1.0 mmol) in 6 mL of CH<sub>2</sub>Cl<sub>2</sub> was added 1-[3-(dimethylamino)-propyl]-3-ethylcarbodiimide hydrochloride (256 mg, 1.32 mmol). The solution was then stirred at rt for 30 min, followed by addition with Example 6 Step 3 amine TFA salt (160 mg, 0.66 mmol) and 4-(dimethylamino)pyridine (244 mg, 2.0 mmol). The solution was then stirred at rt overnight. The mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub> (5 mL) and washed sequentially with H<sub>2</sub>O, 10% citric acid, brine, then dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated to give the title compound (350 mg) which was used without further purification (LC/Mass, + ion): 442 (M+H).

## Step 3



- 5 An oven-dried 10-mL round bottomed flask was charged with Step 2 compound (350 mg, 0.79 mmol), imidazole (108 mg, 1.58 mmol), pyridine (3 mL). The flask under argon was cooled to  $-30^{\circ}\text{C}$ . Slow addition of  $\text{POCl}_3$  (0.30 mL, 3.16 mmol) gave after mixing a thick slurry. The slurry was
- 10 mixed at  $-30^{\circ}\text{C}$  for 3 h and the volatiles evaporated. Dichloromethane (5 mL) was then added and the insoluble solid was removed by filtration. The organic layer was washed with  $\text{H}_2\text{O}$ , 10% citric acid, brine and dried over  $\text{Na}_2\text{SO}_4$ . Removal of solvent gave crude desired nitrile
- 15 (330 mg) (LC/Mass, + ion): 424 (M+H).

## Step 4



- 20 Trifluoroacetic acid (3.3 mL) was added to a stirred solution of Step 3 compound (330 mg, 0.58 mmol) in 3.3 mL  $\text{CH}_2\text{Cl}_2$ . The solution was then stirred at rt for 30 min, a few drops of water were added and the mixture mixture
- 25 stirred for 0.5 h. The mixture was diluted with  $\text{CH}_2\text{Cl}_2$  (5 mL) and concentrated under reduced pressure to a thick oil. The product was purified by reverse phase preparative column chromatography on a YMC S5 ODS 20x100 mm column to give the title compound, 59 mg, 17%.
- 30 Purification conditions: gradient elution from 10% methanol/water/0.1 TFA to 90% methanol/water/ 0.1 TFA

over 15 min; 5 min hold at 90% methanol/water/0.1 TFA.

Flow rate: 20 mL/min. Detection wavelength: 220.


Retention Time 10 Min. (LC/Mass, + ion): 210 (M+H)...

- 5 **General Method B:** Claisen rearrangement sequence to Boc-protected amino acids.



- 10 General method B affords the quaternary Boc-protected amino acids. Examples 30-47 contain the vinyl sidechain by coupling amino acids of which Scheme 4, compound 20 is representative. Cyclopentanone was olefinated under Horner-Emmons conditions to afford 17 which was reduced
- 15 to the allylic alcohol 18 using DIBAL-H in toluene -78 °C to rt. Allylic alcohol 18 was esterified with N-Boc glycine using DCC/DMAP in CH<sub>2</sub>Cl<sub>2</sub> to give 19. Glycine ester 19 was subjected to a Lewis acid mediated Claisen rearrangement by complexation with anhydrous zinc
- 20 chloride and deprotonation at -78°C with lithium diisopropylamide followed by warming to ambient temperature to afford 20.

## Scheme 4, General Method B, Examples 30-47



a. Triethylphosphonoacetate, NaH, THF 0°C to RT b. DIBAL-H, toluene, -78°C to RT c. N-Boc glycine, DCC, DMAP, CH<sub>2</sub>Cl<sub>2</sub>, RT  
d. ZnCl<sub>2</sub>, THF, LDA, -78°C to RT

## Step 1

5 Cyclopentylideneacetic acid ethyl ester.

To a flame-dried 500-mL round-bottomed flask containing NaH (5.10 g of a 60% dispersion in mineral oil, 128 mmol, 1.10 equiv) in 120 mL anhydrous THF at 0°C under argon was added triethylphosphonoacetate (25.6 mL, 128 mmol, 1.10 equiv) dropwise through an addition funnel. The mixture was allowed to warm to rt, stirring for an additional 1 h. A solution of cyclopentanone (10.3 mL, 116 mmol) in 10 mL anhydrous THF was added dropwise over 20 min through an addition funnel, and the mixture was allowed to stir at rt for 2.5 h. Ether (200 mL) and water (100 mL) were then added, and the layers were separated. The organic phase was washed successively with water (100 mL) and brine (100 mL), dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated under reduced pressure, giving 17.5 g (98%) of the desired ester as a colorless oil.




## Step 2

2-Cyclopentylideneethanol.

To a flame-dried 500-mL round-bottomed flask containing  
5 cyclopentylideneacetic acid ethyl ester (17.5 g, 113  
mmol) in 100 mL anhydrous toluene at  $-78^{\circ}\text{C}$  under argon was  
added DIBAL-H (189 mL of a 1.5 M solution in toluene, 284  
mmol, 2.50 equiv) dropwise over a 30 min period through  
an addition funnel, and the mixture was then allowed to  
10 warm to rt, stirring for 18 h. The reaction mixture was  
then recooled to  $-78^{\circ}\text{C}$ , and quenched by the careful  
addition of 30 mL anhydrous MeOH. Upon warming to rt, 1  
N Rochelle's salt (100 mL) was added, and the mixture was  
stirred 90 min. The biphasic reaction mixture was then  
15 diluted with  $\text{Et}_2\text{O}$  (200 mL) in a separatory funnel, and the  
layers were separated. The organic layer was then washed  
with brine (100 mL), dried ( $\text{Na}_2\text{SO}_4$ ), and concentrated  
under reduced pressure. Purification by flash column  
chromatography (silica gel,  $\text{CH}_2\text{Cl}_2$  / EtOAc, 10:1) gave  
20 11.6 g (92%) of the desired allylic alcohol as a  
colorless oil.

## Step 3

25 (2-Cyclopentylideneethyl)-N-(tert-Butyloxycarbonyl)  
glycinate.




To a flame-dried 500-mL round-bottomed flask containing  
30 N-(tert-butyloxycarbonyl)glycine (13.45 g, 76.75 mmol) in  
100 mL  $\text{CH}_2\text{Cl}_2$  at rt was added Step 2 compound (8.61 g,  
76.75 mmol, 1.00 equiv) in 20 mL  $\text{CH}_2\text{Cl}_2$ , followed by  
dicyclohexylcarbodiimide (16.63 g, mmol, 1.05 equiv) in  
80 mL  $\text{CH}_2\text{Cl}_2$ . To this reaction mixture was then added 4-

dimethylaminopyridine (0.94 mg, mmol, 0.10 equiv), and the mixture was allowed to stir overnight. The reaction mixture was then filtered through a medium sintered-glass funnel, rinsing with 100 mL CH<sub>2</sub>Cl<sub>2</sub>, and concentrated under reduced pressure. The crude product was then purified by flash chromatography (silica gel, hexanes/EtOAc, 20:1 to 1:1 gradient) to give 19.43 g (94%) of the desired glyciny ester as a colorless oil.

10 Step 4

N-(tert-Butyloxycarbonyl)(1'-vinylcyclopentyl)-glycine



15

A flame-dried 500-mL round-bottomed flask under argon was charged with ZnCl<sub>2</sub> (11.8 g, mmol, 1.20 equiv) and 20 mL toluene. The mixture was heated under vacuum with vigorous stirring to azeotrope off any traces of moisture with the distilling toluene, repeating this process (2 x). The flask was then cooled to rt under argon, (2-cyclopentylideneethyl) N-(tert-butyloxycarbonyl)glycinate (19.36 g, 71.88 mmol) was added via cannula as a solution in 180 mL THF, and the mixture was then cooled to -78°C.

25 In a separate flame-dried 200-mL round-bottomed flask containing diisopropylamine (26.3 mL, mmol, 2.60 equiv) in 90 mL THF at -78°C was added n-butyllithium (71.89 mL of a 2.5 M solution in hexanes, mmol, 2.5 equiv), and the mixture was allowed to warm to 0°C for 30 min before


30 recooling to -78°C. The lithium diisopropylamine thus generated was then added via cannula to the ZnCl<sub>2</sub> ester mixture dropwise at a steady rate over 40 min, and the

resultant reaction mixture was allowed to slowly warm to rt and stir overnight. The yellow reaction mixture was then poured into a separatory funnel, diluted with 300 mL Et<sub>2</sub>O, and the resultant organic solution was washed successively with 300 mL 1N HCl and 300 mL brine, dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated under reduced pressure. Purification by flash chromatography (silica gel, 3% MeOH in CH<sub>2</sub>Cl<sub>2</sub> with 0.5% HOAc) gave 17.8 g (92%) of the desired amino acid product as a white solid. (FAB MH<sup>+</sup> 270).

10

### Example 30


**General Method C:** Peptide coupling to 4,5-methano-prolinamide, amide dehydration and final deprotection.



15


The TFA salt of amide **13** was coupled to a variety of racemic quaternary protected amino acids using HOBT/EDC in DMF at rt to give a D/L mixture of diastereomers at the N-terminal amino acid. The desired L diastereomer was chromatographically isolated either as the amide **21** or as the nitrile **22**. Nitrile **22** was obtained by treatment of the amide with POCl<sub>3</sub>/imidazole in pyridine at -20°C. The final target **23** was obtained by deprotection under acidic conditions using TFA in CH<sub>2</sub>Cl<sub>2</sub>.

## Scheme 5, General Method C



a. EDAC, HOBT, DMF b.  $\text{POCl}_3$ , pyridine, imidazole,  $-20^\circ\text{C}$  c. TFA,  $\text{CH}_2\text{Cl}_2$ , RT


## Step 1



5


Example 6 Step 3 compound (877 mg, 3.65 mmol) and N-Boc  
cyclopentylvinylamino acid, described in Step 4 of  
10 general method B (1.13 g, 4.20 mmol) were dissolved in 20  
mL anhydrous DMF, cooled to  $0^\circ\text{C}$  and to this mixture was  
added EDAC (1.62 g, 8.4 mmol), HOBT hydrate (2.54 g, 12.6  
mmol, and TEA (1.27 g, 12.6 mmol) and the reaction was  
allowed to warm to rt and stirred for 24 h. The reaction  
15 mixture was taken up in EtOAc (100 mL), washed with  $\text{H}_2\text{O}$  (3  
x 20 mL), dried ( $\text{Na}_2\text{SO}_4$ ), and purified by silica gel flash  
column chromatography (100% EtOAc) to give 1.38 g (86%)  
of Step 1 compound ( $\text{MH}^+$ , 378).

## Step 2



- 5 Step 1 compound (1.38 g, 3.65 mmol) and imidazole (497 mg, 7.30 mmol) were dried by toluene azeotrope (5 mL x 2), dissolved in 10 mL anhydrous pyridine, cooled to -30°C under nitrogen gas and POCl<sub>3</sub> (2.23 g, 14.60 mmol) was added by syringe. The reaction was complete after 1 h
- 10 and was evaporated to dryness and the remainder purified by two sequential flash column chromatographies over silica gel. The first column (100% EtOAc) was used to isolate the mixture of diastereomers (1.15 g, 88%) from the by-products of the reaction. The second column
- 15 (gradient of 25% EtOAc/hexanes to 50% EtOAc/hexanes) was run to resolve the mixture of diastereomers and provided 504 mg of the desired Step 2 nitrile (MH+360).

## Step 3




20

- Step 2 compound (32 mg, 0.09 mmol) was dissolved in 1 mL of CH<sub>2</sub>Cl<sub>2</sub> and 1 mL of TFA was added and the reaction
- 25 stirred for 30 min at rt and was evaporated to dryness. The product was purified by reverse phase preparative column chromatography on a YMC S5 ODS 20 X 250 mm column to give 12 mg of the TFA salt (lyophilized from water or isolated after evaporation of eluent and trituration with
- 30 ether) the title compound. Purification conditions: gradient elution from 10% methanol/water/0.1 TFA to 90%

methanol/water/0.1 TFA over 18 min; 5 min. hold at 90% methanol/water/0.1 trifluoroacetic acid. Flow rate: 20 mL/min. Detection wavelength: 220.







- 5 Examples 30-39 were prepared by the methods outlined in General Method B and General Method C starting from cyclopentanone, cyclobutanone, cyclohexanone, cycloheptanone, cyclooctanone, cis-3,4-dimethylcyclopentanone, and 4-pyranone,
- 10 cyclopropaneethylhemiacetal, acetone, and 3-pentanone respectively.

**Table 2**




15

Example	R	MS [M + H]
30		260
31		246
32		274
33		288

34		302
35		288
36		276
37*		232
38		234
39		262


\* Step 3 compound was prepared by the method described in Tetrahedron Letters 1986, 1281-1284.

#### Example 40



5


Step 1



Step 1 compound was prepared employing general method B starting from cyclopentanone and 2-fluoro-triethylphosphonoacetate instead of triethylphosphonoacetate.


5

Step 2




10 Title compound was prepared by the peptide coupling of Step 1 acid followed by dehydration and final deprotection as described in general method C [MS (M+H) 278].

15

Example 41

Step 1.




20

Step 1 compound was prepared employing general method B starting from cyclobutanone and 2-fluoro-triethylphosphonoacetate instead of triethylphosphonoacetate.

25




## Step 2




5. Title compound was prepared by the peptide coupling of Step 1 acid followed by dehydration and final deprotection as described in general method C. MS (M+H) 264.

10

Example 42

## Step 1




15

Step 1 compound was prepared employing general method B starting from cyclopentanone and triethylphosphono-propionate instead of triethylphosphonoacetate.


20

## Step 2




5 Title compound was prepared by the peptide coupling of  
Step 1 acid followed by dehydration and final  
deprotection as described in general method C. MS (M+H)  
274

10

Example 43

## Step 1




15

Step 1 compound was prepared employing general method B  
starting from cyclobutanone and triethylphosphono-  
propionate instead of triethylphosphonoacetate.

20

## Step 2




5 Title compound was prepared by the peptide coupling of  
 Step 1 acid followed by dehydration and final  
 deprotection as described in general method C. MS (M+H)  
 260.

10

Example 44


**General Method D:** Oxidative cleavage of vinyl  
 substituent by ozonolysis. The protected  
 cyclopentylvinyl nitrile **22** was treated with ozone for 6-  
 15 8 min and subjected to a reductive quench with sodium  
 borohydride to furnish the hydroxymethyl analog **24**  
 directly. This compound was deprotected under acidic  
 conditions with TFA in CH<sub>2</sub>Cl<sub>2</sub> at 0°C to give the target  
 compound **25**.

20

**Scheme 6, General Method D, Examples 44,46,48**


a. O<sub>3</sub>, MeOH:CH<sub>2</sub>Cl<sub>2</sub>, 10:4, -78 C; then NaBH<sub>4</sub>, -78 C to 0 C, 79%  
 b. TFA:CH<sub>2</sub>Cl<sub>2</sub>, 1:2, 0 degrees C.

## Step 1



Cyclopentylvinyl compound prepared in Step 2 of general  
5 method C (1.28 g, 3.60 mmol) was dissolved in 56 mL of a  
2:5 mixture of  $\text{CH}_2\text{Cl}_2$ :methanol, cooled to  $-78^\circ\text{C}$  and was  
treated with a stream of ozone until the reaction mixture  
took on a blue color, at which time,  $\text{NaBH}_4$  (566 mg, 15.0  
mmol, 4.2 equiv) was added and the reaction was warmed to  
10  $0^\circ\text{C}$ . After 30 min, the reaction was quenched with 2 mL  
saturated aqueous  $\text{NaHCO}_3$  and then warmed to rt. The  
reaction mixture was evaporated to dryness and taken up  
in EtOAc. A small amount of water was added to dissolve  
the inorganics and the layers separated. The EtOAc layer  
15 was dried ( $\text{Na}_2\text{SO}_4$ ), filtered and evaporated to an oil that  
was purified by flash column chromatography on silica gel  
with EtOAc to give 922 mg (71%) of Step 1 compound.  
MS(M+H) 364.

## 20 Step 2



Step 1 compound (900 mg, 2.48 mmol) was dissolved in 60  
mL of  $\text{CH}_2\text{Cl}_2$ , cooled to  $0^\circ\text{C}$  and treated with 20 mL of  
25 freshly distilled TFA. The reaction was complete in 80  
min and the mixture was evaporated to dryness and  
purified by preparative HPLC (YMC S5 ODS 30 x 100 mm, 18  
minute gradient 80% Solv A:Solv B to 100% Solv B, Solvent  
A = 10% MeOH-90% $\text{H}_2\text{O}$ -0.1% TFA, Solvent B = 90% MeOH-10%

H<sub>2</sub>O -.1% TFA, collected product from 5.1-6.5 min) to give, after lyophilization from water, 660 mg (71%) of title compound, TFA salt as a white lyophilate. (MH+264).


5

Example 45


**General Method E:** Oxidative cleavage of vinyl substituent by osmium tetroxide-sodium periodate followed by sodium borohydride reduction to alcohol. The cyclobutylolfin **26** was treated with osmium tetroxide and sodium periodate in THF:water, 1:1, and the intermediate aldehyde was isolated crude and immediately reduced with sodium borohydride to give **27** in 56% yield. Standard deprotection conditions using TFA afforded the target compound **28**.

15

**Scheme 7, General Method E, Examples 45, 47**




a. OsO<sub>4</sub>, THF:H<sub>2</sub>O, 1:1; NaIO<sub>4</sub>; workup, then NaBH<sub>4</sub>, MeOH, RT. 56%  
b. TFA:CH<sub>2</sub>Cl<sub>2</sub>, 1:2, 0 degrees C to RT.




20

## Step 1



5 N-Boc protected cyclobutylvinyl compound (Example 31, prepared by general method C) (0.16 g, 0.46 mmol) was dissolved in 10 mL of a 1:1 mixture of THF:water and treated with  $\text{OsO}_4$  (12 mg, catalyst) and  $\text{NaIO}_4$  (0.59 g, 2.76 mmol, 6 equiv). After 2 h, the reaction mixture was  
10 diluted with 50 mL of ether and 10 mL of water. The layers were equilibrated and the organic fraction was washed one time with  $\text{NaHCO}_3$  solution, dried over  $\text{MgSO}_4$  and concentrated to give a dark oil. The oil was diluted with 10 mL of methanol and treated with  $\text{NaBH}_4$  (0.08 g, 2.0  
15 mmol). The mixture turned very dark and after 30 min was diluted with ether and the reaction was quenched with aqueous  $\text{NaHCO}_3$  solution. The mixture was equilibrated and layers separated. The organic fraction was washed with solutions of  $\text{NaHCO}_3$  and 0.1 M  $\text{HCl}$ . The organics were  
20 dried ( $\text{MgSO}_4$ ) and concentrated to give 90 mg (56%) of the Step 1 compound as a dark oil.

## Step 2




25

Step 1 compound (90 mg, 0.26 mmol) was dissolved in 3 mL of  $\text{CH}_2\text{Cl}_2$ , cooled to  $0^\circ\text{C}$  and treated with 3 mL of freshly distilled TFA. The reaction was complete in 80 min and evaporated to dryness and purified by preparative HPLC


- 5 (YMC S5 ODS 30 x 100 mm, 10 minute gradient 100%A to 100%B, Solvent A = 10% MeOH-90% $\text{H}_2\text{O}$ -0.1% TFA, Solvent B = 90% MeOH-10%  $\text{H}_2\text{O}$  -0.1% TFA, to give, after removal of water, 50 mg (60%) of title compound. (MH+250).

10


Table 3



Example	R	Method of Preparation	[M + H]
44		Ozonolysis/ borohydride	264
45		Osmium/periodate/ borohydride	250
46		Ozonolysis/ borohydride	278
47		Osmium/periodate/ borohydride	292
48		Ozonolysis/ borohydride	292

Example 49

## 5 Step 1




Part A. A 50-mL flask was charged with dihydro-4,4-  
10 dimethyl-2,3-furandione (5.0 g, 39.0 mmol), acetic acid  
(10 mL), sodium acetate (3.82 g, 39.0 mmol) and  
hydroxylamine hydrochloride (2.71 g, 39.0 mmol). The  
reaction mixture was stirred for 2 h at rt and  
concentrated under reduced pressure to remove most of the  
15 acetic acid. The remainder was poured into water (100  
mL) and the aqueous phase extracted with EtOAc (3 X 40  
mL). The organics were dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated  
to a colorless oil which solidified on standing.

20 Part B. A 200-mL round bottomed flask was charged with  
Part A solid (@ 39 mmol) and diluted with 80 mL of  
ethanol and 39 mL of 2N HCl (78 mmol). The mixture was  
treated with 1.0 g of 5% Pd/carbon and the mixture  
degassed. The flask was placed under an atmosphere of H<sub>2</sub>  
25 for 8 h. The mixture was filtered through celite and the  
filtrate concentrated to an off white solid.



Part C. A 250-mL round bottomed flask was charged with Part B solid and diluted with THF (50 mL) and water (15 mL). The mixture was treated with di-tert-butylidicarbonate (12.7 g, 117 mmol) and sodium bicarbonate (10.0 g, 117 mmol). After 4 h of stirring the mixture was diluted with 50 mL of ether and 50 mL of water. The layers were separated and the organic fraction dried over  $\text{MgSO}_4$  and concentrated. The residue was purified by flash column chromatography on silica gel with 30% EtOAc in hexanes to give 2.00 g (22% overall) of Step 1 compound as a white solid.


## Step 2



To a stirred solution of Step 1 compound (1.00 g, 3.80 mmol) in THF (20 mL) at rt under nitrogen was added LiOH hydrate (0.16 g, 3.80 mmol) and then water (5 mL). The reaction was stirred at 40°C for 0.5 h and then cooled to rt. The mixture was concentrated to dryness and the remainder was stripped from THF (2X), toluene (2X) and THF (1X). The remaining glass was diluted with 5 mL of THF and treated with imidazole (0.63 g, 9.19 mmol) followed by t-butyl-dimethylsilyl chloride (1.26 g, 8.36 mmol). The reaction was stirred overnight and quenched with 10 mL of methanol. After 1 h of stirring the mixture was concentrated. An additional portion of methanol was added and the mixture concentrated. The oil was diluted with ether and 0.1 N HCl (pH 2). The layers were equilibrated and aqueous drawn off. The organic

fraction was dried over  $\text{MgSO}_4$  and concentrated to give 1.25 g (83%) of Step 2 compound as a colorless glass.

### Step 3



5


The Title compound was prepared by the peptide coupling of Step 2 carboxylic acid with Example 6 Step 3 amine, followed by dehydration and deprotection as outlined in General Method C. MS (M+H) 238.

10

**General Method F:** Catalytic Hydrogenation of vinyl substituent. As shown in Scheme 8, the protected vinyl substituted amino acid **20** was transformed to the corresponding saturated analog **29** by catalytic hydrogenation using 10% Pd/C and hydrogen at atmospheric pressure.

15

### Scheme 8, General Method F, Examples 50-56



a. 10% Pd/C, 1 atm  $\text{H}_2$ , MeOH, 12h, 100%

20

### Step 1.


The N-(tert-Butyloxycarbonyl)(1'-vinylcyclopentyl)glycine (2.23 g, 8.30 mmol) was dissolved in 50 mL MeOH and placed in a hydrogenation vessel purged with argon. To this mixture was added 10% Pd-C (224 mg, 10% w/w) and the reaction stirred under 1 atm  $\text{H}_2$  at rt for 12 h. The reaction was filtered through celite and concentrated and

25

purified by flash column chromatography on silica gel with 1:9 methanol:CH<sub>2</sub>Cl<sub>2</sub> to give the Step 1 compound as a glass. (FAB MH+ 272)


- 5 Examples 50-56 were prepared by the peptide coupling of amino acids (where the vinyl substituent has been hydrogenated according to general method F) followed by dehydration and deprotection as described in general method C.

10

Table 4

Example	R1, R2	MS [M + H]
50	Cyclopentyl	262
51	cyclobutyl	248
52	cycloheptyl	290
53	4-pyranyl	278
54	methyl, methyl	236
55	ethyl, ethyl	264
56	methyl, ethyl	250


15

Example 57

The title compound in Example 57 was prepared by the peptide coupling of the isopropyl cyclobutane amino acid.

(where the olefin substituent has been hydrogenated according to general method F) followed by dehydration and deprotection as described in general method C.

5


Example 58

The title compound in Example 58 was prepared by the peptide coupling of the isopropyl cyclopentane amino acid (where the olefin substituent has been hydrogenated according to general method F) followed by dehydration and deprotection as described in general method C. MS (M+H) 276

15 **General Method G:** L-Amino acids synthesized by Asymmetric Strecker Reaction. Commercially available adamantyl carboxylic acid was esterified either in MeOH with HCl at reflux or using trimethylsilyldiazomethane in Et<sub>2</sub>O/methanol to give **30**. The ester was reduced to the alcohol **31** with LAH in THF and then subjected to a Swern oxidation to give aldehyde **32**. Aldehyde **32** was transformed to **33** under asymmetric Strecker conditions with KCN, NaHSO<sub>3</sub> and R-(-)-2-phenylglycinol. The nitrile of **33** was hydrolyzed under strongly acidic conditions using 12M HCl in HOAc to give **34**. The chiral auxiliary was removed by catalytic reduction using Pearlman's catalyst in acidic methanol under 50 psi hydrogen to give **35** and the resulting amino group was protected as the t-butylcarbamate to give **36**.


30

## Scheme 9, General Method G, Examples 59-64



a. LAH, THF, 0 C to RT, 96% b. ClCOCOCl, DMSO, CH<sub>2</sub>Cl<sub>2</sub>, -78 C, 98% c. R-(-)-2-Phenylglycinol, NaHSO<sub>3</sub>, KCN d. 12M HCl, HOAc, 80 C, 16h, 78 % e. 20% Pd(OH)<sub>2</sub>, 50 psi H<sub>2</sub>, MeOH:HOAc, 5:1 f. (Boc)<sub>2</sub>O, K<sub>2</sub>CO<sub>3</sub>, DMF, 92%, 2 steps

## Step 1



5

Adamantane-1-carboxylic acid (10.0 g, 55 mmol, 1 equiv) was dissolved in a mixture of Et<sub>2</sub>O (160 mL) and MeOH (40 mL), and was treated with trimethylsilyl diazomethane (2.0 M in hexane, 30 mL, 60 mmol, 1.1 equiv) and stirred at rt for 3 h. The volatiles were then removed by rotary evaporation and the product purified by flash column chromatography on silica gel (5x15 cm) with 40% CH<sub>2</sub>Cl<sub>2</sub>/hexanes to give the product as a white crystalline solid (10.7 g, 100%).

15

## Step 2



- Step 1 compound (10.7 g, 0.055 mmol, 1 equiv) was dissolved in anhydrous THF (150 mL) under argon and was treated with a solution of  $\text{LiAlH}_4$  (1 M in THF, 69 mL, 69 mmol, 1.25 equiv). After stirring at rt for 1.5 h, the reaction was cooled to  $0^\circ\text{C}$  and quenched sequentially with  $\text{H}_2\text{O}$  (5.1 mL), 15% aq NaOH (5.1 mL), and  $\text{H}_2\text{O}$  (10.2 mL). After stirring at rt for 15 min, the slurry was vacuum filtered, and the solids washed with EtOAc (2x100 mL). The filtrate was concentrated by rotary evaporation and the resulting solid purified by flash column chromatography on silica gel (5x15 cm) with 10% EtOAc/ $\text{CH}_2\text{Cl}_2$ . This afforded the Step 2 product as a white solid (8.74 g, 96%).


## Step 3



- An oven-dried 3-neck flask equipped with 125-mL addition funnel was charged with anhydrous  $\text{CH}_2\text{Cl}_2$  (150 mL) and anhydrous DMSO (10.3 mL, 0.145 mol, 2.5 equiv) under argon atmosphere and cooled to  $-78^\circ\text{C}$ . Slow dropwise addition of oxalyl chloride (6.7 mL, 0.0768 mol, 1.32 equiv) followed by stirring for 15 min provided an activated DMSO adduct. This was treated with a solution of Step 2 compound (9.67 g, 58.2 mmol, 1 equiv) in dry  $\text{CH}_2\text{Cl}_2$  (75 mL) and the reaction allowed to stir for 1 h. The resulting white mixture was then treated dropwise with triethylamine (40.5 mL, 0.291 mol, 5 equiv). After


30 min, the cooling bath was removed, and the reaction quenched sequentially with cold 20% aq  $\text{KH}_2\text{PO}_4$  (25 mL) and cold  $\text{H}_2\text{O}$  (150 mL). After stirring at rt for 15 min the mixture was diluted with  $\text{Et}_2\text{O}$  (400 mL) and the layers were separated. The organics were washed organic with cold 10% aq  $\text{KH}_2\text{PO}_4$  (3x150 mL) and satd aq  $\text{NaCl}$  (100 mL). The organics were dried ( $\text{Na}_2\text{SO}_4$ ), filtered and concentrated. The residue was purified by flash column chromatography on silica gel (5x10 cm) with  $\text{CH}_2\text{Cl}_2$  to give the Step 3 compound as a white solid (9.40 g, 98%).

## Step 4



Step 3 compound (9.40 g, 57 mmol, 1 equiv) was suspended in  $\text{H}_2\text{O}$  (145 mL) and cooled to  $0^\circ\text{C}$ . The mixture was treated with  $\text{NaHSO}_3$  (5.95 g, 57 mmol, 1 equiv),  $\text{KCN}$  (4.0 g, 59 mmol, 1.04 equiv), and a solution of (R)-(-)-phenylglycinol (8.01 g, 57 mmol, 1 equiv) in  $\text{MeOH}$  (55 mL). The resulting mixture was stirred at rt for 2 h, then refluxed for 16 h. The mixture was cooled to rt, and 200 mL of  $\text{EtOAc}$  added. After mixing for 15 min the layers were separated. The aqueous fraction was extracted with  $\text{EtOAc}$ . The combined  $\text{EtOAc}$  extracts were washed with brine (50 mL), dried over anhydrous  $\text{Na}_2\text{SO}_4$ , filtered and the filtrate concentrated. The product was purified by flash column chromatography on silica gel (6.4x20 cm) with 20%  $\text{EtOAc}$ /hexanes to give the desired (R,S) product as a white solid (11.6 g, 37.4 mmol, 65%): MS m/e 311 ( $\text{M}+\text{H}$ ) $^+$ .


## Step 5



The Step 4 nitrile (5.65 g, 18 mmol) was heated in conc. HCl (120 mL) and HOAc (30 mL) at 80°C for 18 h, at which time the reaction was cooled in an ice bath. Vacuum filtration of the resulting precipitate afforded the desired product as a white solid (5.21 g, 14 mmol, 78%). MS m/e 330 (m+H)<sup>+</sup>.

10


## Step 6



The Step 6 compound (5.21 g, 14 mmol) was dissolved in MeOH (50 mL) and HOAc (10 mL), and hydrogenated with H<sub>2</sub> (50 psi) and Pearlman's catalyst (20% Pd(OH)<sub>2</sub>, 1.04 g, 20% w/w) for 18 h. The reaction was filtered through a PTFE membrane filter and the catalyst washed with MeOH (3x25 mL). The filtrate was concentrated by rotary evaporation to afford a white solid. The product was used in Step 7 without further purification.

20

## Step 7




25

The crude Step 6 compound (@ 14 mmol) was dissolved in anhydrous DMF (50 mL) under argon and treated with K<sub>2</sub>CO<sub>3</sub>




(5.90 g, 42 mmol, 3 equiv) and di-tert-butylidicarbonate (3.14 g, 14 mmol, 1 equiv) under argon at rt. After 19 h, the DMF was removed by rotary evaporation (pump) and the residue dried further under reduced pressure. The residue was mixed with H<sub>2</sub>O (100 mL) and Et<sub>2</sub>O (100 mL), the layers separated, and the alkaline aqueous with Et<sub>2</sub>O (2x100 mL) to remove the by-product from the hydrogenolysis step. The aqueous was cooled to 0°C, diluted with EtOAc (200 mL), and stirred vigorously while carefully acidifying the aqueous to pH 3 with 1N aq HCl. The layers separated and the aqueous extracted with EtOAc (100 mL). The combined EtOAc extracts were washed with brine (50 mL), dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and the filtrate concentrated by rotary evaporation. The residue was purified by SiO<sub>2</sub> flash column (5x12 cm) with 5% MeOH/CH<sub>2</sub>Cl<sub>2</sub> + 0.5% HOAc. The product was chased with hexanes to afford the product as a white foam (4.07 g, 13 mmol, 92%): MS m/e 310 (m+H)<sup>+</sup>.


20

Example 59

The title compound in Example 59 was prepared by the peptide coupling of the Step 7 compound in general method G followed by dehydration and deprotection as described in general method C. MS m/e 300 (m+H)<sup>+</sup>.


Example 60

## Step 1



5  
A solution of KMnO<sub>4</sub> (337 mg, 2.13 mmol, 1.1 equiv) in 2% aq KOH (6 mL) was heated to 60°C and Step 7 compound in general method G (600 mg, 1.94 mmol, 1 equiv) was added in portions, and heating increased to 90°C. After 1.5 h, 10 the reaction was cooled to 0°C, EtOAc (50 mL) was added, and the mixture was carefully acidified to pH 3 with 1N HCl. The layers were separated and the aqueous was extracted with EtOAc (50 mL). The combined organic 15 extracts were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The residue was purified by flash column chromatography on silica gel (3.8x15 cm) with 2% (200 mL), 3% (200 mL), 4% (200 mL), and 5% (500 mL) MeOH/CH<sub>2</sub>Cl<sub>2</sub> + 0.5% HOAc. After isolation of the 20 product, the material was chased with hexanes to afford a white solid (324 mg, 51%): MS m/e 326 (m+H)<sup>+</sup>.


## Step 2



25

The Step 1 compound (404 mg, 1.24 mmol, 1 equiv) was dissolved in anhydrous DMF (10 mL) under argon and cooled to 0°C. The following were added in order: Example 6 Step 3 salt (328 mg, 1.37 mmol, 1.1 equiv), HOBT (520 mg, 3.85 mmol, 3.1 equiv), EDAC (510 mg, 2.61 mmol, 2.1 equiv), and TEA (0.54 mL, 3.85 mmol, 3.1 equiv). The reaction mixture was allowed to warm to rt overnight and the DMF removed by rotary evaporation (pump). The remainder was dried further under vacuum. The residue was dissolved in EtOAc (100 mL), washed with satd aq NaHCO<sub>3</sub> (50 mL) and satd aq NaCl (25 mL), dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated by rotary evaporation. The product was purified flash column chromatography on silica gel (3.8x15 cm) with a gradient of 6% (200 mL), 7% (200 mL), and 8% (500 mL) MeOH/CH<sub>2</sub>Cl<sub>2</sub> to give the product as a white solid (460 mg, 1.06 mmol, 85%): MS m/e 434 (m+H)<sup>+</sup>.

## Step 3




20

The Step 2 compound (95 mg, 0.22 mmol, 1 equiv) was dissolved in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (2.5 mL) under argon and cooled to -78°C. The mixture was treated with diisopropylethylamine (65 µL, 0.37 mmol, 1.7 equiv), and triethylsilyl triflate (75 µL, 0.33 mmol, 1.5 equiv), and stirred at 0°C for 1.5 h. The reaction was mixed with MeOH (0.5 mL), silica gel (200 mg) and H<sub>2</sub>O (2 drops) and stirred at rt for 18 h. The solvent was removed by rotary evaporation and the residue purified flash column

30


chromatography on silica gel (2.5x10 cm) with 4% MeOH/CH<sub>2</sub>Cl<sub>2</sub> to afford the product (92 mg, 0.17 mmol, 77%): MS m/e 548 (m+H)<sup>+</sup>.

5 Step 4



The Step 3 compound (90 mg, 0.16 mmol, 1 equiv) was dissolved in anhydrous pyridine (2 mL) under argon and cooled to -30°C. Treatment with imidazole (24 mg, 0.35 mmol, 2.1 equiv) and phosphorous oxychloride (66 µL, 0.67 mmol, 4.1 equiv), and continued stirring at -30°C for 45 min gave a thick slurry. Volatiles were by rotary evaporation and the cake dried further under reduced pressure. The product was purified by flash column chromatography on silica gel (2.5x10 cm) with 7% EtOAc/CH<sub>2</sub>Cl<sub>2</sub> to afford the product as a white foam (76 mg, 87%): MS m/e 530 (m+H)<sup>+</sup>


20 Step 5




The Step 4 compound (76 mg, 0.14 mmol) was dissolved in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (1 mL) and cooled to 0°C and treated with TFA (1 mL) and H<sub>2</sub>O (2 drops) and stirred for 1.5 hr at 0°C. The solvents were removed by rotary evaporation and the residue was chased with toluene (5 mL) and dried

under reduced pressure. Trituration with Et<sub>2</sub>O afforded the title compound as a white solid (54 mg, 88%): MS m/e 316 (m+H)<sup>+</sup>.

5

Example 61


## Step 1



10

An oven-dried flask purged with argon was charged with anhydrous CH<sub>2</sub>Cl<sub>2</sub> (3 mL) and cooled to -78°C. Treatment with diethylaminosulfur trifluoride (DAST, 60 µL, 0.45 mmol, 1.5 equiv), followed by a solution of the Example 60 Step 2 compound (131 mg, 0.30 mmol, 1 equiv) in dry CH<sub>2</sub>Cl<sub>2</sub> (3 mL). After 15 min, the reaction was poured into a separatory funnel containing satd aq NaHCO<sub>3</sub> (25 mL) and the layers were separated. The aqueous fraction was extracted with CH<sub>2</sub>Cl<sub>2</sub> (25 mL), then the combined organic extracts were washed with brine (10 mL), dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated. The product was purified by flash column chromatography on silica gel (2.5x10 cm) with 5% MeOH/CH<sub>2</sub>Cl<sub>2</sub> to give Step 1 compound (124 mg, 0.29 mmol, 94%): MS m/e 436 (m+H)<sup>+</sup>.


## Step 2



The fluorinated amide from Step 1 (161 mg, 0.37 mmol, 1 equiv) was dissolved in anhydrous pyridine (4 mL) under argon and cooled to  $-30^{\circ}\text{C}$ . The mixture was treated with imidazole (54 mg, 0.77 mmol, 2.1 equiv) and phosphorous oxychloride (143  $\mu\text{L}$ , 1.52 mmol, 4.1 equiv) and stirred at  $-30^{\circ}\text{C}$  for 40 min. The solvent was removed by rotary evaporation and dried further under reduced pressure. The product was purified by flash column chromatography on silica gel (2.5x10 cm) with 5% EtOAc/ $\text{CH}_2\text{Cl}_2$  to give the Step 2 compound as a white foam (126 mg, 82%): MS m/e 418 ( $m+H$ )<sup>+</sup>.


15

## Step 3




The Step 2 compound (125 mg, 0.30 mmol) was dissolved in TFA/ $\text{CH}_2\text{Cl}_2$  (1:1 v/v, 2 mL), and stirred at rt. After 30 min, the solvents were removed by rotary evaporation, the remainder was chased with toluene (2x5 mL), and the solid dried under reduced pressure. Trituration with  $\text{Et}_2\text{O}$  afforded the title compound as a white solid (93 mg, 0.21 mmol, 72%): MS m/e 318 ( $m+H$ )<sup>+</sup>.

25

Example 62


## Step 1




5

The Step 1 compound was prepared beginning with 2-adamantanal and elaborated to the homochiral Boc-amino acid by an asymmetric Strecker synthesis according to  
10 general method G.


## Step 2




15 The title compound in Example 62 was prepared by the peptide coupling of the 2-adamantyl amino acid described in Step 1 followed by dehydration and deprotection as described in general method C. MS (M+H) 300.

Example 63

## Step 1




5 An oven-dried flask equipped with a condenser and drying tube was charged with norbornane-2-carboxylic acid (4.92 g, 35 mmol, 1 equiv) and treated with bromine (2.1 mL, 41 mmol, 1.15 equiv) and phosphorous trichloride (0.153 mL, 1.8 mmol, 0.05 equiv). The mixture was heated at 85°C for 7 h protected from light. Additional bromine (0.4 mL, 7.8 mmol, 0.22 equiv) was added with continued heating for 1 h. The mixture was cooled to rt, and Et<sub>2</sub>O (100 mL) was added. The mixture was washed with 10% aq NaHSO<sub>3</sub> (50 mL), H<sub>2</sub>O (2x50 mL), and brine (25 mL). The ether fraction was dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated by rotary evaporation. The product was purified by flash column chromatography on silica gel (5x15 cm) with 2% to 4% MeOH/CH<sub>2</sub>Cl<sub>2</sub> + 0.5% HOAc. The product was chased with hexanes to remove residual HOAc. The isolated material consists of two inseparable materials (4.7 g), which was used without further purification in the next step.



25 The crude product from above, exo-2- bromonorbornane-1- carboxylic acid (4.7 g, impure) in Et<sub>2</sub>O (80 mL) and MeOH (20 mL), was mixed with trimethylsilyldiazomethane (2.0 M in hexane, 11.8 mL, 23.6 mol), and stirred at rt for 1 h.



Solvent was removed by rotary evaporation, and purification of the oil by flash column chromatography on silica gel (5x18 cm) with a gradient of CH<sub>2</sub>Cl<sub>2</sub>/hexanes (600 mL each of 20% and 30%) followed by CH<sub>2</sub>Cl<sub>2</sub> afforded the product as a white solid (3.97 g, 0.017 mol, 79% for 2 steps): MS m/e 233/235 (m+H)<sup>+</sup>.




Methyl exo-2-bromonorbornane-1-carboxylate (2.0 g, 8.58 mmol, 1 equiv) was dissolved in anhydrous THF (50 mL) in an oven-dried 3-neck flask equipped with a condenser, and purged with argon. The mixture was treated with AIBN (288 mg, 1.71 mmol, 0.2 equiv) and tributyltin hydride (3.6 mL, 12.87 mmol, 1.5 equiv), and then heated to reflux for 2 h. The flask was cooled to rt, and the THF was removed by rotary evaporation to give the crude product. The product was purified by flash column chromatography on silica gel (5x10 cm) with 5% EtOAc/hexanes. The resulting material was used in the next step without further purification.

## Step 2




The Step 1 compound was prepared beginning with 1-norbornyl methyl carboxylate and elaborated to the homochiral Boc amino acid by an asymmetric Strecker synthesis according to general method G.

## Step 3




The title compound in Example 63 was prepared by the peptide coupling of the 1-norbornyl amino acid described in Step 2, followed by dehydration and deprotection as described in general method C. MS (M+H) 260.

Example 64


10

## Step 1



15 The Step 1 compound was prepared beginning with 4-formylpyran and elaborated to the homochiral Boc amino acid by an asymmetric Strecker synthesis according to general method G.

## 20 Step 2




The title compound in Example 64 was prepared by the peptide coupling of the 4-pyranyl amino acid described in Step 2, followed by dehydration and deprotection as described in general method C. MS (M+H) 250.

5

**General Method H:** Strecker Synthesis of Racemic Amino Acids.


**Scheme 10, General Method H, Examples 65-66**

10



a. celite, PCC,  $\text{CH}_2\text{Cl}_2$ , RT, 91% b.  $\text{NH}_4\text{Cl}$ , NaCN, MeOH; 12M HCl, HOAc;  $(\text{Boc})_2\text{O}$ , TEA, DMF.


Step 1



15


To a stirred solution of 1-phenylcyclopentylmethanone (5.00 g, 26.3 mmol) in 25 mL of THF at 0°C was added LAH (52 mL, 52 mmol, 1M) in THF. The reaction mixture was slowly warmed to rt and then refluxed for 18 h. The reaction was quenched according to the Fieser procedure: careful addition of 2 mL of water; 6 mL of 15% NaOH in water; and 2 mL of water. The biphasic mixture was diluted with 100 mL of ether and the granular white solid filtered off. The ether fraction was dried over  $\text{Na}_2\text{SO}_4$  and evaporated to give 4.30 g (93%) of the Step 1 compound.

## Step 2



- 5 To a stirred solution of Step 1 compound (0.80 g, 4.50 mmol) in 15 mL of  $\text{CH}_2\text{Cl}_2$  at rt was added celite (5 g) followed by PCC (1.95 g, 5.00 mmol). After stirring for 3 h the reaction mixture was diluted with 40 mL of  $\text{CH}_2\text{Cl}_2$  and filtered through celite. The filtrate was filtered  
10 an additional time through silica gel resulting in a colorless filtrate. The  $\text{CH}_2\text{Cl}_2$  fraction was evaporated to give 0.72 g (91%) of the aldehyde as a colorless oil.

## Step 3



15


- To a 50-mL round-bottomed flask containing Step 2 compound (0.72 g, 4.20 mmol) in 8 mL of water at rt was added NaCN (0.20 g, 4.20 mmol) followed by  $\text{NH}_4\text{Cl}$  (0.20 g,  
20 5.00 mmol). To this reaction mixture was then added methanol (8 mL) and the mixture was allowed to stir overnight. The reaction mixture was then extracted with ether (2X15 mL), dried ( $\text{MgSO}_4$ ) and concentrated under reduced pressure to give the crude Strecker product.

25


To a 100-mL round-bottomed flask containing the crude Strecker product was added 10 mL of HOAc and 10 mL of conc. HCl. The mixture was refluxed overnight. The mixture was concentrated under reduced pressure to give a

yellow solid. The solid was triturated with 5 mL of 1:1 mixture of ether and hexanes. The white solid was treated with triethylamine (1.4 mL, 9.99 mmol) and di-tert-butylidicarbonate (1.00 g, 4.60 mmol) in 50 mL DMF. After 4 h the pH of the mixture was adjusted to 9 with saturated  $\text{Na}_2\text{CO}_3$  soln. After an additional 3 h of stirring the mixture was extracted with 1:1 ether and hexanes and the aqueous fraction acidified to pH 2 with 5%  $\text{KHSO}_4$  solution. The aqueous phase was washed with ether (2 X 40 mL), the organics dried ( $\text{MgSO}_4$ ), and evaporated to an oil that was purified by silica gel flash chromatography with 8:92 methanol: $\text{CH}_2\text{Cl}_2$  to give 0.3 g (23%) of the Boc-protected amino acid as a light oil (M-H, 318).

15

Example 65

## Step 1




20

The synthesis of the Step 1 compound was described in general method H for the Strecker synthesis of racemic amino acids.


25

## Step 2




The title compound in Example 65 was prepared by the peptide coupling of the cyclopentylphenyl amino acid described in Step 1 and general method H followed by dehydration and deprotection as described in general method C. MS (M+H) 310.

10

Example 66


## Step 1



15

Step 1 compound was prepared using racemic Strecker synthesis according to general method H starting from 2,2-dimethyl-phenylacetic acid.


## 20 Step 2



The title compound in Example 66 was prepared by the peptide coupling of the dimethylphenyl amino acid described in step 1 followed by dehydration and deprotection as described in general method C. MS (M+H)

5 284.

Example 67




10 Step 1

N-(Benzyloxycarbonyl) succinimide (5.6 g, 22.4 mmol) was dissolved in  $\text{CH}_2\text{Cl}_2$  (25 mL) and the solution was added to a cooled ( $0^\circ\text{C}$ ) and stirred solution of diethyl

15 aminomalonate hydrochloride (5.0 g, 23.6 mmol) and triethylamine (13.4 mL, 95 mmol) in  $\text{CH}_2\text{Cl}_2$  (125 mL). The resulting solution was stirred at  $0^\circ\text{C}$  for 10 min and then at rt for 1 h. The solution was washed with 10% citric acid (2 x 50 mL), 10% sodium hydrogen carbonate (2 x 50

20 mL), and water (50 mL) and was then dried ( $\text{Na}_2\text{SO}_4$ ) and evaporated to afford diethyl N-benzyloxycarbonylamino-malonate as a colorless oil, which crystallized upon standing at  $0^\circ\text{C}$  (6.3 g) (LC/Mass + ion):310 (M+H).

25 Step 2



Step 1 compound (6.18 g, 20 mmol) was dissolved in dry ethanol (30 mL) and added to a solution of sodium


30 ethoxide (2.85 g, 8.8 mmol; 21% w/w solution in ethanol (6 mL). A solution of 3-methyl-2-butenal (1.68 g, 20

mmol) in ethanol (12 mL) was added, and the solution stirred at 25°C for 24 h. Acetic acid (0.56 mL) was then added the solution hydrogenated at 50 psi for 24 h using 10% Pd/C (2.0 g) as catalyst. The solution was filtered, 5 evaporated and the residue chromatographed on silica with CH<sub>2</sub>Cl<sub>2</sub> / EtOAc (9:1) to give 2,2-dicarboethoxy-3,3-dimethyl-pyrrolidine (1.6 g) (LC/Mass, + ion): 244 (M+H).

10 This diester (850 mg) was refluxed in 5 M hydrochloric acid (10 mL)/TFA (1 mL) for 8 h to give, after evaporation, a powdery white solid. Crystallization from methanol/ether gave 3,3-dimethyl-dl-proline hydrochloride (190 mg) as white crystals mp 110-112°C.

15


### Step 3



Step 2 compound (173 mg, 0.97 mmol) was dissolved in DMF (3 mL)/ water (3 mL). To this clear solution was added triethylamine (0.46 mL, 3.18 mmol) and di-*t*-butyl dicarbonate (0.23 g, 1.06 mmol), and the reaction mixture was stirred at rt for 5 h. The solution was evaporated and the residue chromatographed on silica column using CH<sub>2</sub>Cl<sub>2</sub>/methanol (9:1) as eluent to yield *t*-butyloxy-carbonyl-3,3-dimethyl-dl-proline (200 mg) as an oil (LC/Mass, + ion): 244 (M+H).




## Step 4




The title compound in Example 67 was prepared by the peptide coupling of the t-butyloxycarbonyl-3,3-dimethyl-dl-proline amino acid described in Step 3 followed by dehydration and deprotection as described in general method C. MS (M+H) 220.

10

Example 68

## Step 1




15

Sodium ethoxide (940 mg of 21 wt% solution in ethanol, 2.9 mmol) in ethanol (2 mL) was added to a stirred solution of diethyl acetamidomalonate (4.31g, 19.8 mmol) in EtOH (23 mL) at rt under argon. The reaction mixture was cooled to 0°C; and trans-2-pentenal (1.51 g, 18.0 mmol) was added dropwise maintaining the reaction temperature at < 5°C. After the addition, the reaction was allowed to warm to rt, stirred for 4 h, then quenched with acetic acid (460 µl). The solution was concentrated in vacuo, and the residue dissolved in EtOAc (25 mL), washed with 10% NaHCO<sub>3</sub> solution (2x5 mL), brine and dried (MgSO<sub>4</sub>). The solution was filtered and concentrated to a


10 mL volume, then heated to reflux and diluted with hexane (20 mL). Upon cooling to rt, the title compound precipitated and was collected to give 3.0 g (50%) of the Step 1 compound (mp 106-109°C; LC/Mass: + ions, 324 M+Na).

## Step 2



10 To a solution of Step 1 compound (2.87 g, 9.5 mmol) and triethylsilane (2.28 mL, 14.3 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (30 mL) under argon was added TFA (7.35 mL, 95.3 mmol) dropwise with stirring while maintaining the internal temperature at 25°C by means of an ice bath. After stirring for 4 h  
15 at rt, the solution was concentrated. The residue was diluted with CH<sub>2</sub>Cl<sub>2</sub> (100 mL), then treated with H<sub>2</sub>O (50 mL) and solid Na<sub>2</sub>CO<sub>3</sub> with vigorous stirring until the mixture was basic. The organic layer was separated, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, then concentrated to give the  
20 Step 2 compound as a yellow oil which was used without further purification (LC/Mass: + ions, 308 M+Na).

## Step 3




25

Step 2 compound (3.73 g, 9.5 mmol) was suspended in 6 N HCl (20 mL) and HOAc (5 mL) and heated at reflux for 20 h. The reaction mixture was then cooled, washed with EtOAc (20 mL), then concentrated to give an oil which  
30 crystallized upon trituration with ether to give the

title compound (1.2 g, 70.6%) (LC/Mass, + ion): 144  
(M+H).


Step 4



5  
Step 3 compound (692 mg, 3.76 mmol) was dissolved in acetone (12 mL)/ water (12 mL). To this clear solution was added triethylamine (1.9 mL, 12.8 mmol) and di-  
10 t-butyl dicarbonate (928 mg, 4.24 mmol). The reaction mixture was stirred at rt for 18 h. The solvents were evaporated and the residue chromatographed on silica with 1:9 methanol:CH<sub>2</sub>Cl<sub>2</sub> to give the Step 4 compound as an oil (LC/Mass: + ions, 266 M+Na).


15

Step 5




Example 68 compound was prepared by peptide coupling of  
20 Step 4 amino acid followed by dehydration and deprotection as described in general method C (MS (M+H) 234).

Example 69



25


## Step 1



Sodium ethoxide (940 mg, 2.9 mmol; 21% w/w solution in  
5 ethanol) in ethanol (2 mL) was added to a stirred  
solution of diethyl acetamidomalonate (4.31 g, 19.8 mmol)  
in EtOH (23 mL) at rt under argon. The reaction mixture  
was cooled to 0°C; and 4-methyl-2-penten-3-ol (1.77 g, 18.0  
mmol) was added dropwise maintaining the reaction  
10 temperature at < 5°C. After the addition, the reaction  
was allowed to warm to rt, stirred for 4 h, then quenched  
with acetic acid (460 µl). The solution was concentrated  
and the remainder dissolved in EtOAc (25 mL). The  
organics were washed with 10% NaHCO<sub>3</sub> solution (2x5 mL),  
15 brine and dried (MgSO<sub>4</sub>). The solution was filtered and  
concentrated to 10 mL volume, then heated to reflux and  
treated with hexane (20 mL). On cooling, the Step 1  
compound precipitated and was collected (3.3 g) (LC/Mass,  
+ ion): 338 (M+Na).

20

## Step 2 .




25 To a solution of Step 1 compound (3.0g, 9.5 mmol) and  
triethylsilane (2.28 mL, 14.3 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (30 mL)  
under argon was added TFA (7.35 mL, 95.3 mmol) dropwise  
with stirring while maintaining the internal temperature  
at 25°C, by means of an ice bath. After stirring for 4 h

at rt, the solution was concentrated, the residue diluted with  $\text{CH}_2\text{Cl}_2$  (100 mL), then treated with  $\text{H}_2\text{O}$  (50 mL) and solid  $\text{Na}_2\text{CO}_3$  with vigorous stirring until the mixture was basic. The organic layer was separated, dried ( $\text{Na}_2\text{SO}_4$ ),  
5 filtered, then concentrated to give the title compound as an oil which was used without further purification (LC/Mass: + ions, 300 M+H).


## Step 3

10




Step 2 compound (3.8 g, 9.5 mmol) was suspended in 6 N HCl (20 mL) and HOAc (5 mL) and heated at reflux for 20  
15 h. The reaction mixture was cooled, washed with EtOAc (20 mL), then concentrated to give an oil which crystallized upon trituration with ether to give the step 3 compound (1.4 g, 76.0%). LC/Mass: + ions, 158 (M+H).

## 20 Step 4




Step 3 compound (728 mg, 3.76 mmol) was dissolved in a  
25 1:1 acetone/water solution (24 mL). To this clear solution was added triethylamine (1.9 mL, 12.8 mmol) and di-*t*-butyl dicarbonate (928 mg, 4.24 mmol). The reaction mixture was stirred at rt for 18 h. The solution was evaporated and the residue chromatographed on silica  
30 column using  $\text{CH}_2\text{Cl}_2$ /methanol (9:1) as eluent to give the title compound as an oil (LC/Mass, + ion): 258 (M+H).

## Step 5




5 Example 69 compound was prepared by peptide coupling of Step 4 amino acid followed by dehydration and deprotection as described in general method C (MS (M+H) 248).

10

Example 70


## Step 1



15

Step 1 compound was prepared by the procedure described in General Method C starting from N-Boc-S-t-butylcysteine.

## Step 2




25


A 25-mL round-bottomed flask equipped with a magnetic stirring bar and N<sub>2</sub> inlet was charged with Step 1 compound

(78 mg, 0.21 mmol) and chloroform (3 mL). The mixture was cooled to 0°C and treated with *m*-chloroperoxybenzoic acid (85 mg, 0.44 mmol) in CHCl<sub>3</sub> (2 mL). After 3 h the solution was diluted with CHCl<sub>3</sub> (7 mL), washed with 5% NaHCO<sub>3</sub> (2x5 mL), H<sub>2</sub>O and dried over Na<sub>2</sub>SO<sub>4</sub>. Removal of solvent gave crude sulfoxide (100 mg), which was used without further purification (LC/Mass, + ions): 384 (M+H).

## 10 Step 3




15 Trifluoroacetic acid (1.5 mL) was added to a cooled (0°C) solution of Step 2 compound (100 mg, 0.26 mmol) in 5 mL CH<sub>2</sub>Cl<sub>2</sub>. The solution was then stirred at 0°C for 1.5 h, diluted with CH<sub>2</sub>Cl<sub>2</sub> (5 mL) and concentrated under reduced pressure to a thick oil. The product was purified by  
20 reverse phase preparative column chromatography on a YMC S5 ODS 20x100 mm column to give the title compound of Example 70, 17 mg, 16%. Purification conditions: gradient elution from 10% methanol/water/0.1 TFA to 90% methanol/water/ 0.1 TFA over 15 min 5 min hold at 90%  
25 methanol/water/0.1 TFA. Flow rate: 20 mL/min. Detection wavelength: 220. Retention Time 10 Min (LC/Mass, + ion): 284 (M+H).

Example 71

30


## Step 1



5 A 25-mL round-bottomed flask equipped with a magnetic stirring bar and N<sub>2</sub> inlet was charged with compound from Example 70, Step 1 (78 mg, 0.21 mmol) in chloroform (3 mL). The mixture was cooled to 0°C and treated with *m*-chloroperoxybenzoic acid (144 mg, 0.84 mmol) in CHCl<sub>3</sub> (2 mL). After 30 min at rt, the solution was diluted with CHCl<sub>3</sub> (7 mL), washed with 5% NaHCO<sub>3</sub> (2x10 mL), H<sub>2</sub>O and dried over Na<sub>2</sub>SO<sub>4</sub>. Removal of solvent gave the crude sulfone (100 mg), which was used without further purification (LC/Mass, + ion): 344 (M+H-Bu).

15

## Step 2



20 Trifluoroacetic acid (1.5 mL) was added to a cooled (0°C) and stirred solution of Step 1 compound (100 mg, 0.26 mmol) in 5 mL CH<sub>2</sub>Cl<sub>2</sub>. The solution was stirred at 0°C for 30 min, diluted with CH<sub>2</sub>Cl<sub>2</sub> (5 mL) and concentrated under reduced pressure to a thick oil. The product was


25 purified by reverse phase preparative column chromatography on a YMC S5 ODS 20x100 mm column to give the title compound, 14 mg, 17%. Purification conditions: gradient elution from 10% methanol/water/0.1 TFA to 90% methanol/water/ 0.1 TFA over 15 min. 5 min hold at 90%

30 methanol/water/0.1 TFA. Flow rate:20 mL/min. Detection



wavelength: 220. Retention Time 10 Min. (LC/Mass, +  
ion): 300 (M+H).


Example 72



The title compound was prepared following a published  
procedure (Sasaki et al, Tetrahedron Lett. 1995, 36,  
10 3149, Sasaki et al. Tetrahedron 1994, 50, 7093) used to  
synthesize (2S,3R,4S)-N-Boc-3,4-methano-L-proline  
carboxylate. The corresponding amide was prepared by  
general method A and deprotected with TFA to give the TFA  
salt also as described in general method A.


15

Example 73




The title compound was prepared by coupling (2S,3R,4S)-  
3,4-methano-L-proline carboxamide-N-trifluoroacetate  
described in Example 72 with L-cyclohexylglycine and then  
dehydrated to the amide with POCl<sub>3</sub>/imidazole and  
deprotected (N-terminal nitrogen) with TFA using general  
25 C (FAB MH+ 248).

Example 74



The title compound was prepared by coupling (2S,3R,4S)-3,4-methano-L-proline carboxamide-N-trifluoroacetate described in Example 72 with L-tert-butylglycine and then  
5 dehydrated to the amide with POCl<sub>3</sub>/imidazole and deprotected (N-terminal nitrogen) with TFA using general C (FAB MH+ 222).


Example 75



10


The title compound was prepared by coupling (2S,3R,4S)-3,4-methano-L-proline carboxamide-N-trifluoroacetate  
15 described in Example 72 with L-valine and then dehydrated to the amide with POCl<sub>3</sub>/imidazole and deprotected (N-terminal nitrogen) with TFA using general C (FAB MH+ 207).

Example 76




20

The title compound was prepared by coupling (2S,3R,4S)-  
25 3,4-methano-L-proline carboxamide-N-trifluoroacetate described in Example 72 with N-(tert-butyloxycarbonyl)-(1'ethylcyclopentyl)glycine described in General Method B and then dehydrated to the amide with POCl<sub>3</sub>/imidazole and deprotected (N-terminal nitrogen) with TFA using general  
30 C (FAB MH+ 262).

Example 77


5 The title compound was prepared by coupling (2S,3R,4S)-  
3,4-methano-L-proline carboxamide-N-trifluoroacetate  
described in Example 72 with N-(tert-butyloxycarbonyl)-  
(1'-vinylcyclopentyl)glycine described in General Method B  
and then dehydrated to the amide with POCl<sub>3</sub>/imidazole and  
10 deprotected (N-terminal nitrogen) with TFA using General  
Method C (FAB MH+ 260).

Example 78

15


N-[(S)-cyclopentylvinyl]-N-tert-butoxycarbonylglycinyll]-  
(2S,4S,5S)-2-cyano-4,5-methano-L-prolylamide (70 mg, 0.19  
mmol) described in General Method C, Step 2 was dissolved  
20 in a mixture of 2 mL t-BuOH / 3 mL THF and N-  
methylmorpholine-N-oxide (33mg, 0.28 mmol) was added  
followed by osmium tetroxide (0.1 mmol, 50 mol%). The  
reaction was quenched with 1 mL of 10% aqueous Na<sub>2</sub>SO<sub>3</sub> and  
was taken up in EtOAc and washed with H<sub>2</sub>O 5 mL, dried  
25 (Na<sub>2</sub>SO<sub>4</sub>), filtered, evaporated and purified by silica gel  
flash chromatography (5% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) to give 41 mg (55%)  
of the protected diol as an oil. The title compound was  
obtained by deprotection of the amine functionality with  
TFA according to General Method C (FAB MH+ 294).

30

Example 79


5 **General Procedure I: Synthesis of Quaternary Amino Acids**  
 Via Michael Addition to Malonates followed by Selective  
 Hydrolysis and Curtius Rearrangement. Examples 79-84.

Cyclohexanone and diethylmalonate underwent Knoevenagel  
 10 condensation mediated by titanium tetrachloride in THF  
 and CCl<sub>4</sub> to give **40**. Copper (I) mediated Grignard  
 addition of methylmagnesium bromide gave **41** which was  
 selectively saponified to **42**. Curtius rearrangement with  
 trapping by benzyl alcohol gave **43** which was converted to  
 15 **44** by a standard deprotection-protection protocol. Ester  
**44** was saponified to give the quaternary amino acid **45**.

**Scheme 11, General Method I**

a. THF, CCl<sub>4</sub>, TiCl<sub>4</sub>, diethylmalonate, 0 °C; pyridine, THF, 0 to RT 72 h b. MeMgBr,  
 CuI, Et<sub>2</sub>O, 0 °C c. 1N NaOH, EtOH, RT 6 days d. Ph<sub>2</sub>PON<sub>3</sub>, TEA, RT to reflux to RT,  
 20 BrOH e. 10% Pd(OH)<sub>2</sub>/C, EtOAc; (Boc)<sub>2</sub>O, K<sub>2</sub>CO<sub>3</sub>, THF f. 1N NaOH, dioxane

## Step 1




5

According to literature procedure (Tetrahedron 1973, 29, 435), a mixture of dry tetrahydrofuran (400 mL) and dry carbon tetrachloride (50 mL) was cooled to 0°C (ice-salt bath) and treated with titanium tetrachloride (22.0 mL, 0.2 mole). The resulting yellow suspension was stirred at 0°C for 5 min, treated sequentially with cyclohexanone (10.3 mL, 0.1 mole) and distilled diethylmalonate (15.2 mL, 0.1 mole) then stirred at 0°C for 30 min. The reaction mixture was then treated with a solution of dry pyridine (32 mL, 0.40 mole) in dry THF (60 mL), stirred at 0°C for 1.0 h, then at rt for 72 h. The reaction mixture was quenched with water (100 mL), stirred for 5 min then extracted with ether (2 x 200 mL). The combined organic extracts were washed with saturated sodium chloride (100 mL), saturated sodium bicarbonate (100 mL) and brine (100 mL), dried over anhydrous magnesium sulfate, filtered and concentrated. Flash chromatography using 5% EtOAc in hexane gave step 1 compound as a light yellow oil. Yield: 5.25 g (22%). MS (M + Na) 263.


25

## Step 2



- 5 According to literature (Org. Syn. VI, 442, 1988; Liebigs Ann. Chem. 1981, 748) a mixture of 3.0 M methylmagnesium iodide (3.1 mL, 9.36 mmol) and cuprous chloride (9.0 mg) was stirred at 0°C (ice-salt water bath), treated with a solution of Step 1 compound (1.5 g, 6.24 mmol) in dry  
10 ether (1.8 mL) over 5 min and stirred at 0°C for 1 h, then at rt for 40 min. The mixture was slowly added to a slurry of ice and water (15 mL), treated dropwise with 10% HCl (3.7 mL) then extracted with EtOAc (3 x 25 mL). The combined organic extracts were washed with 1% sodium  
15 thiosulfate (2.0 mL) and saturated sodium chloride (2.0 mL), dried over anhydrous magnesium sulfate, filtered, and concentrated. Flash chromatography on a silica gel column using 5% ether in hexane (1.0 L) gave step 2 compound as a clear syrup. Yield: 1.09 g, (68%). MS  
20 (M+H) 257.

## Step 3




25

A solution of Step 2 compound (1.09 g, 4.03 mmol) in a mixture of methanol (5.4 mL) and water (2.7 mL) was treated with 1N sodium hydroxide (4.84 mL, 4.84 mmol or

1.2 equiv) and stirred at rt for 6 days. The reaction mixture still showed the presence of starting material, so THF (4.0 mL) was added and the entire mixture stirred for another 2 days. The solution was evaporated to dryness and the resulting syrup partitioned between water (8.0 mL) and ether (15 mL). The aqueous phase was acidified with 1N hydrochloric acid (4.8 mL) to pH 2-3 and extracted with EtOAc (3 x 25 mL). The combined organic extracts were washed with brine (10.0 mL), dried over anhydrous magnesium sulfate, filtered, and concentrated to give step 3 compound as a thick syrup. Yield: 875 mg, (95.1%). MS (M + H) 229.

Or alternately: solutions of the diester in a mixture of ethanol, THF, dioxane and water or mixtures thereof may be hydrolyzed with sodium hydroxide.

#### Step 4




20

According to literature (J. Org. Chem 1994, 59, 8215), a solution of Step 3 compound (0.875 g, 3.83 mmol) in dry benzene (4.0 mL) was treated with triethylamine (0.52 mL, 3.83 mmol) and diphenylphosphoryl azide (0.85 mL, 3.83 mmol), refluxed under nitrogen for 1 h and cooled to rt. The solution was treated with benzyl alcohol (0.60 mL, 5.75 mmol or 1.5 equiv), refluxed for 17 h, cooled then diluted with ether (40 mL). The solution was washed with 10% aqueous citric acid (2x3 mL), back-extracting the citric acid wash with ether (40 mL). The combined

30

organic extracts were washed with 5% sodium bicarbonate (2x3 mL), dried (MgSO<sub>4</sub>), filtered, and concentrated. Flash chromatography on silica gel of the crude product with 10% EtOAc in hexane (1.0 L) gave step 4 compound as  
5 a clear thick syrup. Yield: 1.15 g (90%). MS(M+H) 334.

### Step 5



10

A solution of Step 4 compound (1.15 g, 3.46 mmol) in EtOAc (60 mL) was treated with palladium hydroxide on carbon (298 mg) and hydrogenated at rt for 20 h. The mixture was filtered through a celite pad and then washing the pad well with EtOAc (3 x 25 mL) then the filtrate was concentrated to give the free amine. A solution of the amine in tetrahydrofuran (12 mL) and water (12 mL) was treated with di-*t*-butyl dicarbonate (1.0 g, 4.58 mmol or 1.48 equiv) and potassium carbonate (854 mg, 6.18 mmol or 2.0 equiv), then stirred at rt for 20 h. The reaction mixture was partitioned between water (8 mL) and diethyl ether (3 x 40 mL) and the combined organic extracts were washed with brine (8 mL), dried ( $\text{MgSO}_4$ ), filtered, and concentrated. Flash chromatography of the crude product with 10% EtOAc in hexane (1 L) gave step 5 compound as a clear thick syrup. Yield: 1.18 g (100%). MS: (M+H) 300.

Other methods can also be employed, for example:

30 According to Tetrahedron Lett. 1988, 29, 2983, where a solution of the benzylcarbamate in ethanol may be treated




with triethylsilane (2 equiv), di-*t*-butyldicarbonate (1.1 equiv), catalytic palladium acetate and triethylamine (0.3 equiv) to give the BOC-protected amine in a "one-pot" manner.

5

Or alternately: Solutions of the benzylcarbamate in methanol may be subjected to hydrogenolysis in the presence of di-*t*-butyldicarbonate to give the BOC-protected amine in a "one-pot" manner.


10

#### Step 6



- 15 A solution of Step 5 compound (1.18 g, 3.09 mmol) in dioxane (8.0 mL) was treated with 1N sodium hydroxide (9.1 mL, 9.1 mmol or 3.0 equiv) and stirred at 60°C (oil bath) for 28 h. The reaction mixture was concentrated to a syrup which was dissolved in water (15 mL) and
- 20 extracted with ether (25 mL). The aqueous phase was acidified to pH 2-3 with 1N hydrochloric acid (9.2 mL) then extracted with EtOAc (3 x 50 mL). The combined organic extracts were washed with saturated sodium chloride (10 mL), dried (MgSO<sub>4</sub>), filtered, and
- 25 concentrated to give Step 6 compound as an off-white solid. Yield: 808 mg (96%). MS (M+H) 272.

## Step 7




5 The title compound was prepared from Step 6 compound according to the procedure in General Method C where the amino acid was coupled, the amide was dehydrated, and the protecting group removed to give the title compound. MS (M+H) 262.

10


Compounds 90-100 were prepared by General Method I and General Method C starting from cyclohexanone, cyclopentanone and cyclobutanone, and employing methyl-, ethyl-, allyl- and propylmagnesium halides as Grignard

15 reagents.


Table 5

Example #	Cycloalkane	R	MS Data M+H
79	cyclohexane	Methyl	262
80	cyclohexane	Ethyl	276
81	cyclopentane	Methyl	248
82	cyclopentane	Allyl	274
83	cyclopentane	Propyl	276
84	cyclobutane	Methyl	234

20

Example 85

---


Step 1

5

According to Example 79: A mixture of dry carbon tetrachloride (50 mL) was cooled to 0°C (ice-salt bath) and treated with titanium tetrachloride (11.0 mL, 0.1 mol). The resulting yellow suspension was stirred at 0°C for 5 min, treated sequentially with cyclopentanone (4.42 mL, 0.05 mol) and distilled diethylmalonate (7.6 mL, 0.05 mol) then stirred at 0°C for 30 min. The reaction mixture was then treated with a solution of dry pyridine (16 mL, 0.20 mol) in dry THF (30 mL), stirred at 0°C for 1.0 h, then at rt for 20 h. The reaction mixture was quenched with water (50 mL), stirred for 5 min then extracted with ether (2 x 100 mL). The combined organic extracts were washed with saturated sodium chloride (50 mL), saturated sodium bicarbonate (50 mL) and brine (50 mL), dried (MgSO<sub>4</sub>), filtered and concentrated. Flash chromatography using 5% EtOAc in hexane gave Step 1 compound as a light yellow oil. Yield: 7.67 g (68%). MS (M + H) 226.


25

## Step 2



- 5 A solution of Step 1 compound (1.00 g, 4.42 mmol) in methanol (50 mL) was treated with 10% Pd/C (0.20 g, 10 mol%) and hydrogenated (balloon pressure) at rt for 20 h. The mixture was diluted with methanol and filtered through a pad of celite. The filtrate was concentrated and purified by flash column chromatography on silica gel with 7% EtOAc in hexanes to give 0.84 g (91%) of Step 2 compound. MS (M+H) 229.


## Step 3



15

- The Step 3 compound was prepared by the process outlined in General Method H, where the ester underwent hydrolysis, Curtius Rearrangement, protecting group exchange, and again final ester hydrolysis.


## Step 4



25

The title compound was prepared from Step 3 compound according to the procedure in General Method C where the amino acid was coupled, the amide was dehydrated, and the protecting group removed to give the title compound. MS (M+H) 234.


Examples 86 and 87 were prepared by the procedures used for Example 85 starting from cyclohexanone and cyclobutanone respectively.



10


Example #	Cycloalkane	Mass Spec
		M+H
85	cyclopentyl	234
86	cyclohexyl	248
87	cyclobutyl	220

### Example 89




15

Step 1



Step 1 compound was prepared in Example 6 Step 1.

Step 2




5

The title compound was prepared from Step 1 compound according to General Method C, where the carboxylic acid  
10 underwent a peptide coupling, the amide dehydration and protecting group removal. MS (M+H) 218.

Examples 90 to 99


Examples of compounds where X = H include the following  
 compounds which may be prepared employing procedures as  
 5 described hereinbefore.



Ex. #	n	x	y	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>
90	0	0	1	t-Bu	H	H	-
91	0	0	1	adamantyl	H	H	-
92	0	0	1		H	H	-
93	0	0	1		H	Me	-
94	0	1	0	t-Bu	H	H	-
95	0	1	0	adamantyl	H	H	-
96	0	1	0		H	H	-
97	0	1	0		H	Me	-
98	1	0	1	H	H	H	t-Bu
99	1	1	0	Me	H	H	t-Bu

Examples 100 to 109

Examples of compounds where  $n = 1$  include the following compounds which may be prepared employing procedures as described hereinbefore.




Ex. #	X	x	y	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>
100	CN	0	1	H	H	H	t-Bu
101	CN	0	1	H	H	H	adamantyl
102	CN	0	1	H	Me	H	
103	CN	0	1		H	Me	H
104	CN	1	0	t-Bu	H	H	H
105	CN	1	0	adamantyl	H	H	Me
106	CN	1	0		Et	H	H
107	CN	1	0	H	H	Me	
108	H	0	1	t-Bu	H	H	H
109	H	1	0	Me	H	H	t-Bu



What is Claimed is:

1. A compound having the structure



5

herein  $x$  is 0 or 1 and  $y$  is 0 or 1, provided that

$x = 1$  when  $y = 0$  and

$x = 0$  when  $y = 1$ ; and wherein

$n$  is 0 or 1;

10

$X$  is H or CN;

$R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are the same or different and are

independently selected from hydrogen, alkyl, alkenyl,

alkynyl, cycloalkyl, cycloalkylalkyl, bicycloalkyl,

tricycloalkyl, alkylcycloalkyl, hydroxyalkyl,

15 hydroxyalkylcycloalkyl, hydroxycycloalkyl,

hydroxybicycloalkyl, hydroxytricycloalkyl,

bicycloalkylalkyl, alkylthioalkyl, arylalkylthioalkyl,

cycloalkenyl, aryl, aralkyl, heteroaryl, heteroarylalkyl,

cycloheteroalkyl or cycloheteroalkylalkyl; all optionally

20 substituted through available carbon atoms with 1, 2, 3,

4 or 5 groups selected from hydrogen, halo, alkyl,

polyhaloalkyl, alkoxy, haloalkoxy, polyhaloalkoxy,

alkoxycarbonyl, alkenyl, alkynyl, cycloalkyl,

cycloalkylalkyl, polycycloalkyl, heteroarylamino,

25 arylamino, cycloheteroalkyl, cycloheteroalkylalkyl,

hydroxy, hydroxyalkyl, nitro, cyano, amino, substituted

amino, alkylamino, dialkylamino, thiol, alkylthio,

alkylcarbonyl, acyl, alkoxycarbonyl, aminocarbonyl,

alkynylaminocarbonyl, alkylaminocarbonyl,

30 alkenylaminocarbonyl, alkylcarbonyloxy,


alkylcarbonylamino, arylcarbonylamino,

alkylsulfonylamino, alkylaminocarbonylamino,


alkoxycarbonylamino, alkylsulfonyl, aminosulfinyl,

aminosulfonyl, alkylsulfinyl, sulfonamido or sulfonyl;

- and  $R^1$  and  $R^3$  may optionally be taken together to form  $-(CR^5R^6)_m-$  where  $m$  is 2 to 6, and  $R^5$  and  $R^6$  are the same or different and are independently selected from hydroxy, alkoxy, H, alkyl, alkenyl, alkynyl, cycloalkyl, halo, amino, substituted amino, cycloalkylalkyl, cycloalkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl, alkylcarbonylamino, arylcarbonylamino, alkoxy carbonylamino, aryloxy carbonylamino, alkoxy carbonyl, aryloxy carbonyl, or alkylaminocarbonylamino, or  $R^1$  and  $R^4$  may optionally be taken together to form  $-(CR^7R^8)_p-$  wherein  $p$  is 2 to 6, and  $R^7$  and  $R^8$  are the same or different and are independently selected from hydroxy, alkoxy, cyano, H, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, cycloalkenyl, halo, amino, substituted amino, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl, alkylcarbonylamino, arylcarbonylamino, alkoxy carbonylamino, aryloxy carbonylamino, alkoxy carbonyl, aryloxy carbonyl, or alkylaminocarbonylamino, or optionally  $R^1$  and  $R^3$  together with




from a 5 to 7 membered ring containing a total of 2 to 4 heteroatoms selected from N, O, S, SO, or SO<sub>2</sub>;




- or optionally  $R^1$  and  $R^3$  together with form a 4 to 8 membered cycloheteroalkyl ring wherein the cycloheteroalkyl ring has an optional aryl ring fused thereto or an optional 3 to 7 membered cycloalkyl ring fused thereto;
- including all stereoisomers thereof;
- and a pharmaceutically acceptable salt thereof, or a prodrug ester thereof, and all stereoisomers thereof.


2. The compound as defined in Claim 1 having the structure:




5 3. The compound as defined in Claim 1 having the structure:



10 4. The compound as defined in Claim 1 having the structure:




15 5. The compound as defined in Claim 1 having the structure:




20 6. The compound as defined in Claim 1 wherein:  
 $R^3$  is H,  $R^1$  is H, alkyl, cycloalkyl, bicycloalkyl, tricycloalkyl, alkylcycloalkyl, hydroxyalkyl, hydroxyalkylcycloalkyl, hydroxycycloalkyl, hydroxybicycloalkyl, or hydroxytricycloalkyl,  
 $R^2$  is H or alkyl, n is 0,  
 25 X is CN.

7. The compound as defined in Claim 1 wherein the cyclopropyl fused to the pyrrolidine has the configuration:



5

8. The compound as defined in Claim 1 having the structure:




15

or a pharmaceutically acceptable salt thereof.

9. The compound as defined in Claim 8 wherein the pharmaceutically acceptable salt is the hydrochloride salt or the trifluoroacetic acid salt.

20


10. The compound as defined in Claim 1 which is



A

(1S, 2(2S), 3S, 5S)

wherein R<sup>1</sup> is alkyl, cycloalkyl, bicycloalkyl,  
 5 tricycloalkyl, alkylcycloalkyl, hydroxyalkyl,  
 hydroxycycloalkyl, hydroxyalkylcycloalkyl,  
 hydroxybicycloalkyl, or hydroxytricycloalkyl,  
 or



B

(1R, 2S, 3(2S), 5S)

10

wherein R<sup>1</sup> is alkyl, cycloalkyl, bicycloalkyl,  
 tricycloalkyl, alkylcycloalkyl, hydroxyalkyl,  
 hydroxycycloalkyl, hydroxyalkylcycloalkyl,  
 hydroxybicycloalkyl, or hydroxytricycloalkyl.

15

11. A pharmaceutical composition comprising a  
 compound as defined in Claim 1 and a pharmaceutically  
 acceptable carrier therefor.

20

12. A pharmaceutical combination comprising a DP4  
 inhibitor compound as defined in Claim 1 and an  
 antidiabetic agent other than a DP4 inhibitor for  
 treating diabetes and related diseases, an anti-obesity  
 agent and/or a lipid-modulating agent.

25

13. The pharmaceutical combination as defined in  
 Claim 12 comprising said DP4 inhibitor compound and an  
 antidiabetic agent.

14. The combination as defined in Claim 13 wherein the antidiabetic agent is 1, 2, 3 or more of a biguanide, a sulfonyl urea, a glucosidase inhibitor, a PPAR  $\gamma$  agonist, a PPAR  $\alpha/\gamma$  dual agonist, an SGLT2 inhibitor, an  $\alpha$ P2 inhibitor, a glycogen phosphorylase inhibitor, an AGE inhibitor, an insulin sensitizer, a glucagon-like peptide-1 (GLP-1) or mimetic thereof, insulin and/or a meglitinide.

10

15. The combination as defined in Claim 14 wherein the antidiabetic agent is 1, 2, 3 or more of metformin, glyburide, glimepiride, glipyrider, glipizide, chlorpropamide, gliclazide, acarbose, miglitol, pioglitazone, troglitazone, rosiglitazone, insulin, Gl-262570, isaglitazone, JTT-501, NN-2344, L895645, YM-440, R-119702, AJ9677, repaglinide, nateglinide, KAD1129, AR-HO39242, GW-409544, KRP297, AC2993, Exendin-4, LY307161, NN2211, and/or LY315902.

20

16. The combination as defined in Claim 13 wherein the compound is present in a weight ratio to the antidiabetic agent within the range from about 0.01 to about 100:1.

25

17. The combination as defined in Claim 12 wherein the anti-obesity agent is a beta 3 adrenergic agonist, a lipase inhibitor, a serotonin (and dopamine) reuptake inhibitor, a thyroid receptor beta compound, an anorectic agent, and/or a fatty acid oxidation upregulator.

30

18. The combination as defined in Claim 17 wherein the anti-obesity agent is orlistat, ATL-962, AJ9677, L750355, CP331648, sibutramine, topiramate, axokine, dexamphetamine, phentermine, phenylpropanolamine, famoxin, and/or mazindol.

35

19. The combination as defined in Claim 12 wherein the lipid modulating agent is an MTP inhibitor, an HMG CoA reductase inhibitor, a squalene synthetase inhibitor, a fibric acid derivative, an upregulator of LDL receptor activity, a lipoxxygenase inhibitor, an ACAT inhibitor, a cholesteryl ester transfer protein inhibitor, or an ATP citrate lyase inhibitor.

20. The combination as defined in Claim 19 wherein the lipid modulating agent is pravastatin, lovastatin, simvastatin, atorvastatin, cerivastatin, fluvastatin, nisvastatin, visastatin, fenofibrate, gemfibrozil, clofibrate, implitapide, CP-529,414, avasimibe, TS-962, MD-700, and/or LY295427.

15

21. The combination as defined in Claim 19 wherein the DP4 inhibitor is present in a weight ratio to the lipid-modulating agent within the range from about 0.01 to about 100:1.

20

22. A pharmaceutical combination comprising a DP4 inhibitor compound as defined in Claim 1 and an agent for treating infertility, an agent for treating polycystic ovary syndrome, an agent for treating a growth disorder and/or frailty, an anti-arthritis agent, an agent for preventing inhibiting allograft rejection in transplantation, an agent for treating autoimmune disease, an anti-AIDS agent, an agent for treating inflammatory bowel disease/syndrome, an agent for treating anorexia nervosa, an anti-osteoporosis agent and/or an anti-obesity agent.

23. A method for treating diabetes, insulin resistance, hyperglycemia, hyperisulinemia, or elevated blood levels of free fatty acids or glycerol, obesity, Syndrome X, dysmetabolic syndrome, diabetic complications, hypertriglyceridemia,

hyperinsulinemia, atherosclerosis, impaired glucose homeostasis, impaired glucose tolerance, infertility, polycystic ovary syndrome, growth disorders, frailty, arthritis, allograft rejection in transplantation, 5 autoimmune diseases, AIDS, intestinal diseases, inflammatory bowel syndrome, nervosa, osteoporosis, or an immunomodulatory disease or a chronic inflammatory bowel disease, which comprises administering to a mammalian species in need of treatment a therapeutically effective 10 amount of a compound as defined in Claim 1.

24. The method as defined in Claim 23 for treating type II diabetes and/or obesity.

15



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau




(43) International Publication Date  
20 September 2001 (20.09.2001)

PCT

(10) International Publication Number  
WO 01/68603 A3

- (51) International Patent Classification<sup>7</sup>: C07D 209/52. A61K 31/403, A61P 3/04, 3/06, 3/10
- (21) International Application Number: PCT/US01/07151
- (22) International Filing Date: 5 March 2001 (05.03.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/188,555 10 March 2000 (10.03.2000) US
- (71) Applicant (for all designated States except US): BRISTOL-MYERS SQUIBB CO. [US/US]; P.O. Box 4000, Lawrenceville-Princeton Road, Princeton, NJ 08543 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): ROBL, Jeffrey, A. [US/US]; 7 Tulip Drive, Newtown, PA 18940 (US). SULSKY, Richard, B. [US/US]; 2 Coventry Square, West Trenton, NJ 08628 (US). AUGERI, David, J. [US/US]; 107 Carter Road, Princeton, NJ 08540 (US). MAGNIN, David, R. [US/US]; 40 Cottage Court, Hamilton, NJ 08690 (US). HAMANN, Lawrence, G. [US/US]; 24 East Riding Drive, Cherry Hill, NJ 08003 (US). BETEBENNER, David, A. [US/US]; 3 Easton Court, Lawrenceville, NJ 08648 (US).
- (54) Title: CYCLOPROPYL-FUSED PYRROLIDINE-BASED INHIBITORS OF DIPEPTIDYL IV, PROCESSES FOR THEIR PREPARATION, AND THEIR USE
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— with international search report
- (88) Date of publication of the international search report:  
14 February 2002
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/68603 A3



(57) Abstract: Dipeptidyl peptidase IV (DP 4) inhibiting compounds are provided having the formula (I) where x is 0 or 1 and y is 0 or 1 (provided that x = 1 when y = 0 and x = 0 when y = 1); n is 0 or 1; X is H or CN; and wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are as described herein. A method is also provided for treating diabetes and related diseases, especially Type II

diabetes, and other diseases as set out herein, employing such DP 4 inhibitor or a combination of such DP 4 inhibitor and one or more of another antidiabetic agent such as metformin, glyburide, troglitazone, pioglitazone, rosiglitazone and/or insulin and/or one or more of a hypolipidemic agent and/or anti-obesity agent and/or other therapeutic agent.

PC7/US 01/07151

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C07D209/52 A61K31/403 A61P3/04 A61P3/06 A61P3/10

#### B. FIELDS SEARCHED

IPC 7 C07D

EPO-Internal, CHEM ABS Data

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
------------	--	-----------------------

1-7, 11,  
23, 24

1-7, 11,  
23, 24

1, 11-13,  
22-24

— / —

☒ Patent family members are listed in annex.

\*8\* document member of the same patent family

23/10/2001

Hass, C

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/07151

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4 379 785 A (WEYER RUDI ET AL) 12 April 1983 (1983-04-12) cited in the application claims 1,5,6 ---	1,11,23, 24
A	EP 0 219 782 A (HOECHST AG) 29 April 1987 (1987-04-29) page 1, formula (I); page 3, substructure K; page 16, line 33 to page 17, line 6 claims 1,2 ---	1,11,23
A	DE 33 24 263 A (HOECHST AG) 17 January 1985 (1985-01-17) claims 1,19,20 ---	1,11
A	DE 39 26 606 A (HOECHST AG) 14 February 1991 (1991-02-14) page 2, formula (I); page 3, substructure (K); page 6, lines 62-66 ---	1
A	US 4 254 057 A (DAY JANET A ET AL) 3 March 1981 (1981-03-03) column 2, line 26 - line 38 ---	1
A	EP 0 007 652 A (SHELL INT RESEARCH) 6 February 1980 (1980-02-06) claim 1 ---	1
A	HANESSION S ET AL: "Probing the Importance of Spatial and Conformational Domains in Captopril Analogs for Angiotensin Converting Enzyme Activity" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, OXFORD, GB, vol. 8, no. 16, 18 August 1998 (1998-08-18), pages 2123-2128, XP004137231 ISSN: 0960-894X the whole document ---	1
A	TVEREZOVSKY V V ET AL: "Synthesis of (2S, 3R, 4S)-3,4-Methanoproline and Analogues by Cyclopropylidene Insertion" TETRAHEDRON, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 53, no. 43, 27 October 1997 (1997-10-27), pages 14773-14792, XP004106307 ISSN: 0040-4020 the whole document ---	1
	-/--	

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/07151

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	SAGNARD I ET AL: "Enantioselective Synthesis of Cyclopropane alpha-Amino Acids: Synthesis of N-Boc-cis-(2S,3R,4S)-3,4-Methanoproline and N-Boc-(2S,3R,4S)-3,4-Methanoglutamic Acid" TETRAHEDRON LETTERS, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 36, no. 18, 1 May 1995 (1995-05-01), pages 3149-3152, XP004028212 ISSN: 0040-4039 the whole document	1
A	WO 99 26659 A (HOOVER DENNIS JAY; TREADWAY JUDITH LEE (US); HULIN BERNARD (US); P) 3 June 1999 (1999-06-03) cited in the application claims 1,6,7	12-14, 22-24
A	US 5 998 463 A (HULIN BERNARD ET AL) 7 December 1999 (1999-12-07) cited in the application claims 1,22,23	12-14, 22,23

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/07151

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 6011155	A	04-01-2000	US 6124305 A	26-09-2000
WO 9947545	A	23-09-1999	AU 3098699 A	11-10-1999
			BG 104863 A	30-04-2001
			BR 9909660 A	21-11-2000
			EP 1064298 A2	03-01-2001
			NO 20004546 A	09-11-2000
			PL 343611 A1	27-08-2001
			WO 9947545 A2	23-09-1999
WO 9967279	A	29-12-1999	DE 19828114 A1	27-01-2000
			AU 4777299 A	10-01-2000
			BR 9911415 A	20-03-2001
			CN 1306541 T	01-08-2001
			WO 9967279 A1	29-12-1999
			EP 1090030 A1	11-04-2001
			NO 20006483 A	19-12-2000
			US 2001020006 A1	06-09-2001
US 4379785	A	12-04-1983	DE 2951135 A1	25-06-1981
			AR 231131 A1	28-09-1984
			AR 230989 A1	31-08-1984
			AR 240922 A1	27-03-1991
			AT 6934 T	15-04-1984
			AU 538129 B2	02-08-1984
			AU 6552380 A	25-06-1981
			CA 1167033 A1	08-05-1984
			DE 3067390 D1	10-05-1984
			DK 540280 A ,B,	20-06-1981
			EP 0031058 A1	01-07-1981
			ES 497661 D0	01-11-1981
			ES 8200665 A1	01-02-1982
			ES 498224 D0	01-02-1982
			ES 8202541 A1	01-05-1982
			ES 498225 D0	01-03-1982
			ES 8202798 A1	16-05-1982
			ES 498226 D0	01-03-1982
			ES 8202799 A1	16-05-1982
			ES 498227 D0	01-02-1982
			ES 8202542 A1	01-05-1982
			ES 498228 D0	01-02-1982
			ES 8202543 A1	01-05-1982
			FI 803937 A ,B,	20-06-1981
			GR 72534 A1	17-11-1983
			HU 184943 B	28-11-1984
			IE 50635 B1	28-05-1986
			IL 61733 A	31-05-1985
			JP 1039424 B	21-08-1989
			JP 1554161 C	04-04-1990
			JP 56108762 A	28-08-1981
			LU 90244 A9	20-07-1998
			MX 8254 A	01-04-1993
			MX 174496 B	19-05-1994
			MX 8352 A	01-11-1993
			MX 6926 E	01-12-1986
			NO 803855 A ,B,	22-06-1981
			PT 72226 A ,B	01-01-1981
			ZA 8007911 A	30-12-1981

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PC1/US 01/07151

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0219782	A	29-04-1987	DE 3536687 A1	16-04-1987
			AT 95064 T	15-10-1993
			AU 594711 B2	15-03-1990
			AU 6389086 A	16-04-1987
			CA 1320904 A1	03-08-1993
			DE 3689099 D1	04-11-1993
			DK 490486 A	16-04-1987
			EP 0219782 A2	29-04-1987
			ES 2059301 T3	16-11-1994
			IE 60767 B	10-08-1994
			JP 62087524 A	22-04-1987
			KR 9308094 B1	25-08-1993
			PT 83535 A ,B	01-11-1986
			US 5231080 A	27-07-1993
			ZA 8607771 A	27-05-1987
DE 3324263	A	17-01-1985	DE 3324263 A1	17-01-1985
			AT 53203 T	15-06-1990
			AU 573227 B2	02-06-1988
			AU 3029884 A	10-01-1985
			CA 1263000 A1	14-11-1989
			CA 1267902 A2	17-04-1990
			DE 3482385 D1	05-07-1990
			DK 330284 A	07-01-1985
			EP 0131226 A2	16-01-1985
			ES 534001 D0	16-04-1985
			ES 8504705 A1	16-07-1985
			ES 535452 D0	16-05-1985
			ES 8505341 A1	01-09-1985
			ES 535453 D0	16-05-1985
			ES 8505342 A1	01-09-1985
			FI 842691 A	07-01-1985
			GR 82242 A1	13-12-1984
			HU 209413 B	30-05-1994
			HU 39160 A2	28-08-1986
			JP 1985439 C	25-10-1995
			JP 7010879 B	08-02-1995
			JP 60051199 A	22-03-1985
			KR 9107884 B1	04-10-1991
			PT 78848 A ,B	01-08-1984
			US 4591598 A	27-05-1986
			ZA 8405160 A	27-02-1985
DE 3926606	A	14-02-1991	DE 3926606 A1	14-02-1991
			AT 94409 T	15-10-1993
			AU 631914 B2	10-12-1992
			AU 6092090 A	14-02-1991
			CA 2023089 A1	12-02-1991
			CS 9003958 A3	12-08-1992
			DD 297063 A5	02-01-1992
			DE 59002721 D1	21-10-1993
			DK 417473 T3	13-12-1993
			EP 0417473 A1	20-03-1991
			ES 2059931 T3	16-11-1994
			HU 54504 A2	28-03-1991
			IE 902910 A1	27-02-1991
			IL 95327 A	31-10-1995
			JP 3083957 A	09-04-1991

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PC1/US 01/07151

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 3926606	A	KR 185969 B1 NO 903532 A NZ 234836 A PL 286442 A1 PT 94957 A ,B US 5231083 A ZA 9006327 A	01-05-1999 12-02-1991 26-05-1997 09-03-1992 22-05-1991 27-07-1993 29-05-1991
US 4254057	A	03-03-1981 AU 522785 B2 AU 4835079 A BR 7903987 A CA 1117127 A1 DE 2962088 D1 DK 266679 A EP 0007652 A1 ES 481872 A1 JP 55007278 A US 4255334 A US 4279821 A ZA 7903152 A	24-06-1982 03-01-1980 11-03-1980 26-01-1982 18-03-1982 28-12-1979 06-02-1980 01-03-1980 19-01-1980 10-03-1981 21-07-1981 25-06-1980
EP 0007652	A	06-02-1980 AU 522785 B2 AU 4835079 A BR 7903987 A CA 1117127 A1 DE 2962088 D1 DK 266679 A EP 0007652 A1 ES 481872 A1 JP 55007278 A US 4255334 A US 4254057 A US 4279821 A ZA 7903152 A	24-06-1982 03-01-1980 11-03-1980 26-01-1982 18-03-1982 28-12-1979 06-02-1980 01-03-1980 19-01-1980 10-03-1981 03-03-1981 21-07-1981 25-06-1980
WO 9926659	A	03-06-1999 AP 911 A AT 205403 T AU 733304 B2 AU 9555898 A BG 104435 A BR 9814698 A CN 1279617 T EP 1032424 A1 HR 20000327 A1 HU 0100272 A2 WO 9926659 A1 NO 20002164 A PL 340643 A1	07-12-2000 15-09-2001 10-05-2001 15-06-1999 31-01-2001 03-10-2000 10-01-2001 06-09-2000 28-02-2001 28-06-2001 03-06-1999 19-07-2000 12-02-2001
US 5998463	A	07-12-1999 NONE	